Supplementary Appendix 1. Questionnaire form.

KNOWLEDGE ASSESSMENT OF DOG BITE AND RABIES MANAGEMENT IN REGIÃO ADMINISTRATIVA ESPECIAL DE OÉ-CUSSE AMBENO (RAEOA)

1. Full Name: …….

2. Age: …….

3. Sex: ☐ Male ☐ Female

4. Profession: ☐ Doctor (medical) ☐ Nurse ☐ Midwife ☐ Pharmacist
☐ Public Health ☐ Allied technician ☐ Student

5. Workplace: ☐ Regional Health Service ☐ Oé-Cusse Referral Hospital ☐ Passabe Community Health Center (CHC)
☐ Quimanteco CHC ☐ Baqui CHC ☐ Oe-Silo CHC ☐ Baocnana CHC

6. What is rabies? ☐ A vaccine-preventable zoonotic disease that affects the central nervous system. ☐ Cardiovascular disease
☐ A disease transmitted by mosquitoes ☐ Mental illness due to stress

7. 99% of cases of rabies virus transmission to humans come from: ☐ Mosquitoes ☐ Snakes ☐ Dogs ☐ Cats

8. The fatality rate for confirmed human rabies is ☐ 25% ☐ 50% ☐ 75% ☐ 100%

9. The rabies virus (rabies) is transmitted to humans through:
☐ Wind ☐ Saliva, through direct contact with mucosa (such as eyes, mouth, or an open wound).
☐ Water ☐ Sunlight

10. Symptoms and signs of rabies in humans include:
☐ Vomiting blood, fainting, headache ☐ Diarrhea, fever, stomach ache
☐ Hydrophobia, photophobia, stiffness, fever ☐ Coughing, sneezing, body aches

11. How many categories of dog-bite wounds are there? ☐ 1 ☐ 3 ☐ 6

12. A category III wound less than 5 cm, not affecting arteries or tendons, does not need stitches! ☐ True ☐ False

13. A dog-bite wound needs to be washed with soap or detergent with running water for 15 minutes! ☐ True ☐ False

14. A patient comes with a wide, deep, open wound, bleeding a lot, and torn skin on their right leg. What category does this fall into?
☐ I ☐ II ☐ III

15. Choose the correct one
☐ The patient comes with a category III dog-bite wound: treat completely and send home
☐ Patient comes with a category III dog-bite wound: suture the wound, give antibiotics, send home
☐ Patient comes with a category III dog-bite wound: clean the wound for 15 minutes, give anti-tetanus vaccine, give anti-rabies vaccine (dose I), RIG, Report to the Surveillance Responsible

16. The anti-rabies vaccine RABIVAX-S 1 dose-1 mL-is administered via:
☐ Intravenous, left arm
☐ Intramuscular, deltoid (>2 years to adult) or antero-lateral muscle (for children <2 years)
☐ All are wrong

17. A 1 mL intramuscular dose of anti-rabies vaccine can be given to everyone, from small children to adults to pregnant and breastfeeding mothers! ☐ True ☐ False

18. A 3-dose anti-rabies vaccine (days 0, 7, and 21) is given to people at risk of being bitten by dogs, such as veterinarians, immigration personnel, and people in the border area with Indonesia. This falls into the category:
☐ Pre-exposure prophylaxis (PrEP) for rabies ☐ Post-exposure prophylaxis (PEP)

19. A 5-dose anti-rabies vaccine (days 0, 3, 7, 14, and 28) given to people bitten by a dog falls into the category:
☐ Pre-exposure prophylaxis (PrEP) for rabies ☐ Post-exposure prophylaxis (PEP)

20. Human Rabies Immunoglobulin (HRIG), dose 20 IU/kg, will be given to people who are bitten by a dog with a wound category:
☐ Category I ☐ Category II and has already received a complete vaccine ☐ Category III and has received vaccine 1st dose <7 days

21. Human Rabies Immunoglobulin (HRIG) will be injected intramuscularly in the: ☐ Buttocks ☐ Leg ☐ Around the wound (infiltrated)

22. Dog-bite cases must be reported to the Municipal Surveillance Responsible, then reported to National Surveillance.
☐ within 24 hours ☐ Within one week ☐ Within one month

23. What is your opinion regarding the socialization and training on dog-bite management and anti-rabies vaccine from INSP-TL in RAEOA?
☐ Not good ☐ Fair ☐ Good ☐ Very good ☐ Excellent