

The THAI *Journal of* SURGERY

Official Publication of the Royal College of Surgeons of Thailand

Vol. 31

October - December 2010

No. 4

Review Article

Epidemiology and Risk Factors of Esophageal Cancer

Somkiat Sunpaweravong, MD

Department of Surgery, Faculty of Medicine, Prince of Songkla University, Songkla, Thailand

Abstract

Esophageal cancer is the eighth most common malignancy in the world. In general, squamous cell carcinoma is more common than adenocarcinoma, except in the USA. The etiology of the cancer is multifactorial, involving such things as smoking, alcohol, dietary habits, viral infections, GERD, environmental carcinogens, and socioeconomic status. This review also shows factors that may protect against the development of esophageal cancer such as a diet rich in fruits and vegetables or consumption of NSAIDs or helicobacter pylori infection in adenocarcinoma.

Key words: Epidemiology, Esophageal Cancer, Risk Factor

Esophageal cancer, the 8th most frequent cancer and the 6th most common cause of cancer death in the world,^{1,2} has a greatly diverse occurrence rate in different regions around the globe. For example, in Linxian, China the occurrence rate of esophageal cancer is 100 per 100,000,³ while the average rate of the whole country of China is 13 per 100,000.^{4,5} Northern Iran, South Africa and southern Thailand are examples of regions in which a higher occurrence rate of esophageal cancer has been found. In Thailand, esophageal cancer is found more often in the south

when compared with other regions of the country. Thailand statistics on the occurrence rate of cancer show that esophageal cancer was found at 8.1 per 100,000 in its male population and only 1.8 in females in Songkla Province in the south. In other regions of Thailand, such as Chiangmai Province in northern Thailand, the occurrence rate of esophageal cancer is 2.5 per 100,000 in males and 0.7 in females; and 1.8 and 0.6 per 100,000 people in the male and female populations respectively of Khonkaen Province in the northeast.⁶

Correspondence address : Somkiat Sunpaweravong, MD, Department of Surgery, Faculty of Medicine, Prince of Songkla University, Songkla, Thailand; Telephone: +66 7445 1401; Fax: +66 7442 9384; E-mail: susomkia@medicine.psu.ac.th

Concerning specific sub-types, esophageal squamous cell carcinoma (ESCC) is generally found more frequently than esophageal adenocarcinoma (EAC) in Asia and Africa, while EAC has a higher frequency of occurrence than ESCC in the US and Europe. This is because the occurrence of gastro-esophageal reflux disease (GERD), which is related to the occurrence of Barrett's esophagus and EAC, is higher in populations in the US and Europe than in other groups. Presently, the occurrence of EAC is increasing while ESCC is gradually decreasing. Since 1990 EAC has been reported as a higher frequency of occurrence than ESCC in the United States.⁷

It is vital to differentiate between ESCC and EAC cancers because of the different risk factors between them. These risk factors concern both consumption behaviors such as cigarette smoking, alcohol drinking, general dietary habits including the use of medications, and other environmental factors such as passive intake of carcinogens, infections and so on.

Tobacco Smoking

Tobacco ingestion is one of the main risk factors in esophageal cancer. Several case-controlled studies have indicated that the amount and length of time of tobacco smoking has a significant relationship with the occurrence of esophageal cancer.⁸⁻¹¹ The occurrence rate is 4.2 times higher in current smokers and 3.4 times in former smokers than nonsmokers. It has also been found that the length of time smoking has more effect on the occurrence of esophageal cancer than does the amount of tobacco smoked.¹² Tobacco smoking is more associated with EAC cancer than ESCC. However, the amount and length of time in tobacco smoking was found to cause more occurrence of esophageal cancer¹³⁻¹⁵ at a rate of 1.5 to 4 times when compared with those who do not smoke.¹⁶⁻¹⁸

Alcohol

Although alcohol ingestion has been found to have less effect on the occurrence of ESCC than tobacco smoking,^{8,10} a study conducted in Taiwan indicated that former alcohol drinkers had a 5.5 times higher risk of ESCC, and current drinkers had a risk rate of 7.6 occurrence rate than nondrinkers. The amount of alcohol ingestion had a greater influence on the occurrence of ESCC than did the length of time.¹² However, no significant relationship was found

between alcohol ingestion and EAC cancer^{13,14,19}. Although no significant occurrence rate of esophageal cancer was found among those who were both drinkers and smokers,²⁰ a higher relative risk has been found among those who were both heavy smokers and drinkers.^{8,12}

Dietary

Certain types of dietary intakes such as meat, pickles and salted fish have been found in some case-controlled studies to be a relative risk for ESCC. It is of interest to note that such foods likely contain N-nitroso compounds or substrates of nitrate and amine.²¹⁻²³ Diets rich in vegetables and fruits, on the other hand, are known to help prevent the occurrence of cancer. Some studies have suggested that the higher intake of vitamin C in such diets might play a role in fighting the impact of N-nitroso compounds in the human body.²⁴⁻²⁶ Some nutritional deficiencies such as selenium and zinc have also been found to be risk factors for esophageal cancer.²⁷⁻³⁰

Higher ingestion of vegetables and fruits was also found to reduce the risk of EAC. This might be due to the antioxidants in vitamins C and E and beta-carotene in those diets.³¹⁻³³ A study conducted with a large number of subjects (521,457) in European countries found that diets containing a larger amount of meat ingestion can increase the risk of getting EAC.³⁴

Nonsteroidal Anti-inflammatory Drugs (NSAIDs)

A 2003 meta-analysis study found that aspirin and other NSAID drugs are associated with EAC rates up to 40% less than non-NSAID users, depending on the amount of drug used,³⁵ and other studies have had similar findings.³⁶⁻³⁹ It has been hypothesized that NSAID use might reduce inflammations which can lead to cancers, although in one study conducted in Linxian, China, in which celecoxib was randomly given to cases, no reduction of esophageal squamous dysplasia was found.⁴⁰

Human Papilloma Virus (HPV) Infection

It is widely known that HPV, particularly HPV16 and 18, which is the main causes of cervical cancer, including certain types of cancers in such areas as the vulva and penis.^{41,42} HPV was reported as being of relative risk to esophageal cancer in only cases using polymerase chain reaction (PCR)⁴³ while there was no

relationship between these two variables in other techniques.⁴⁴⁻⁴⁷ It might be possible that the finding of HPV in some esophageal cancer patients using the PCR technique was due to some sort of contamination. To date, the International Agency for Research on Cancer (IARC) has concluded that there is no proven evidence to state that HPV causes esophageal cancer.⁴⁸

Epstein-Barr Virus (EBV)

It is widely known that EBV is one of the risk factors for nasopharyngeal carcinoma.⁴⁹ It has also been found in some studies to have a relationship with the occurrence of esophageal cancer.^{50,51} All studies finding the relationships between EBV and esophageal cancer, however, have been done using the PCR technique in which contamination of EBV can occur in lymphocytes of cancer tissues. In contrast when ISH (in situ hybridization) or PCR in cell lines techniques, which are free from lymphocyte contamination, were used, no significant relationship between the virus and esophageal cancer was found.⁵¹⁻⁵⁸

Helicobacter Pylori

H. pylori, a major cause of stomach cancer,^{59,60} was not found to have any correlation with the occurrence of EAC. Three meta-analysis studies reported that *H. pylori* found in the stomach reduced up to 50 percent possibilities of EAC.⁶¹⁻⁶³ It has been hypothesized that *H. pylori* reduces the gastric acid and therefore reduces Gastro-Esophageal Reflux Disease (GERD) which is the cause of Barrett's esophagus and EAC.⁶⁴ There is no known relationship between ESCC and *H. pylori*.

N-Nitroso Compounds

N-Nitroso compounds contain in animal samples have been found to be a relative risk for cancer in the nasal cavity, esophagus, and stomach.⁶⁵⁻⁶⁸ To date there is no clear explanation on how these N-Nitroso compounds are responsible for cancers in human beings. These compounds can be taken into our body by cigarette smoking, ingestion of certain food such as pickled fish, pork sausages, salted beef, salted fish, and also from endogenous synthesis within the human body.^{69,70} No biomarkers were found to detect physical changes in the human body, to date.

N-Nitroso compounds comprised of Nitrosamines and Nitrosamides are the result of a biochemical synthesis of nitrites and amines or amides. Nitrites are

compounds resulting from synthesis of nitrates found in foods by certain types of bacteria in the oral cavity.^{71,72}

Meta-analysis studies in esophageal cancer have shown a high level of correlation between nitrosamines and certain types of foods that contain nitrites and nitrosamines such as animal meat.⁷³

Polycyclic Aromatic Hydrocarbons (PAH)

PAHs are compounds resulting from the incomplete combustion of organic substances such as grilled animal meat and other substances which contribute to environmental air pollution such as cigarette smoke or smoke from burning coal.⁷⁴⁻⁷⁶ PAHs have been recognized as a risk factor for several types of cancer including some types of skin, lungs, and urinary bladder cancers.⁷⁷ Cigarette smoke, which has a high PAH content, is known to be highly correlated to esophageal cancer. However, as with N-nitroso compounds, there is yet no known biomarker to correlate changes in the human body with the intake of these compounds. There are, therefore, no empirical explanations for the relationships between these compounds and esophageal cancers. Chinese patients in Linxian who were found to have a high occurrence of ESCC were also found to have large quantities of urinary PAH markers reflect exposure only in the 24 to 72 hours before urine collection.⁷⁸⁻⁸¹

Acetaldehyde

Acetaldehyde can be taken into the human body in several ways, but the most prevalent source is alcohol. When taken into the human body, the enzyme alcohol dehydrogenase (ADH) transforms ethanol into acetaldehyde which is further transformed into acetate by acetaldehyde dehydrogenase (ALDH). Laboratory studies show that acetaldehyde causes point of mutation in human lymphocytes, cellular proliferation, and inhibits DNA repair. Acetaldehyde is also a risk factor for esophageal cancer.⁸² However, the IARC has said that more empirical studies are needed before acetaldehyde is accepted as a known cause of esophageal cancer.⁸³

Gastro-Esophageal Reflux Disease (GERD)

GERD is widely accepted as one of the risk factors of EAC. Lagergren et al. reported a close relationship between GERD and EAC.⁸⁴ In general the occurrence rate of EAC is 8 times higher in people with GERD than

those without, while the occurrence rate has risen to 20 times in cases of acute GERD patients.⁸⁵⁻⁸⁷ GERD changes the esophageal membrane into Barrett's esophagus and this may finally develop into EAC. There is a risk rate of 0.5-1 per cent a year for the occurrence of EAC in Barrett's esophagus patients.⁸⁸⁻⁹⁰ However, GERD and Barrett's esophagus are not found to have any relationship to ESCC.

Obesity

A high Body Mass Index (BMI) is one of the risk factors for EAC.⁹⁰⁻⁹³ A meta-analysis study reported a positive relationship between EAC and overweight (BMI = 25-30 kg/m²) and obese individuals (BMI >30 kg/m²), at levels of 2 to 3 times respectively above the norm.^{90,91} These findings corresponded with studies among Americans and Europeans which found that overweight and obese people were more susceptible to EAC. This might be because obesity increases abdominal pressures and therefore increases the occurrence rate of GERD⁹⁴ which can later develop into Barrett's esophagus and EAC, although another study also reported a higher occurrence rate of EAC in overweight people even without GERD.⁹⁵

Socioeconomic Status

Socioeconomic status is determined by earnings, education and jobs. People with lower socioeconomic status have been found to have a higher risk of esophageal cancer than those with a higher socioeconomic status. Study conducted in Sweden⁹⁶ has reported an apparent relationship between the occurrence of both ESCC and EAC and socio-economic status in which ESCC had a higher level of relationship.⁹⁷

REFERENCE

1. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;55:74-108.
2. Kamanger F, Dores GM, Anderson WF. Patterns of cancer incidence, mortality and prevalence across five continents: defining priorities to reduce cancer disparities in different geographic regions of the world. *J Clin Oncol* 2006;24:2137-50.
3. Ke L. Mortality and incidence trends from esophagus cancer in selected geographic areas of China circa 1970-90. *Int J Cancer* 2002;102:271-4.
4. Wang Z, Tang L, Sun G, et al. Etiological study of esophageal squamous cell carcinoma in an endemic region: a population-based case control study in Huaian, China. *BMC Cancer* 2006;6:287.
5. Tran GD, Sun XD, Abnet CC, et al. Prospective study of risk factors for esophageal and gastric cancers in the Linxian general population trial cohort in China. *Int J Cancer* 2005;113:456-63.
6. Sriplung H, Wiangnon S, Sontipong S, et al. Cancer incidence trends in Thailand, 1989-2000. *Asian Pac J Cancer Prev* 2006;7:239-44.
7. Holmes RS, Vaughan TL. Epidemiology and pathogenesis of esophageal cancer. *Semin Radiat Oncol* 2007;17:2-9.
8. Hashibe M, Boffetta P, Janout V, et al. Esophageal cancer in Central and Eastern Europe: tobacco and alcohol. *Int J Cancer* 2007;120:1518-22.
9. Jiang JM, Zeng XJ, Chen JS, et al. Smoking and mortality from esophageal cancer in China: a large case-control study of 19,734 male esophageal cancer deaths and 104,846 living spouse controls. *Int J Cancer* 2006;119:1427-32.
10. Wu IC, Lu CY, Kuo FC, et al. Interaction between cigarette, alcohol and betel nut use on esophageal cancer risk in Taiwan. *Eur J Clin Invest* 2006;36:236-41.
11. Wu M, Zhao JK, Hu XS, et al. Association of smoking, alcohol drinking and dietary factors with esophageal cancer in high- and low-risk areas of Jiangsu Province, China. *World J Gastroenterol* 2006;12:1686-93.
12. Lee CH, Lee JM, Wu DC, et al. Independent and combined effects of alcohol intake, tobacco smoking and betel quid chewing on the risk of esophageal cancer in Taiwan. *Int J Cancer* 2005;113:475-82.
13. de Jonge PJ, Steyerberg EW, Kuipers EJ, et al. Risk factors for the development of esophageal adenocarcinoma in Barrett's esophagus. *Am J Gastroenterol* 2006;101:1421-9.
14. Ranka S, Gee JM, Johnson IT, et al. Non-steroidal anti-inflammatory drugs, lower oesophageal sphincter-relaxing drugs and oesophageal cancer. A case-control study. *Digestion* 2006;74:109-15.
15. Whiteman DC, Sadeghi S, Pandeya N, et al. Combined effects of obesity, acid reflux and smoking on the risk of adenocarcinomas of the oesophagus. *Gut* 2008;57:173-80.
16. Crew KD, Neugut AI. Epidemiology of upper gastrointestinal malignancies. *Semin Oncol* 2004;31:450-64.
17. Lagergren J. Adenocarcinoma of oesophagus: what exactly is the size of the problem and who is at risk? *Gut* 2005;54 (Suppl 1):i1-5.
18. Pera M, Manterola C, Vidal O, Grande L. Epidemiology of esophageal adenocarcinoma. *J Surg Oncol* 2005;92:151-9.
19. Freedman ND, Abnet CC, Leitzmann MF, et al. A prospective study of tobacco, alcohol, and the risk of esophageal and gastric cancer subtypes. *Am J Epidemiol* 2007;165:1424-33.
20. Brown LM, Hoover R, Silverman D, et al. Excess incidence of squamous cell esophageal cancer among US Black men: role of social class and other risk factors. *Am J Epidemiol* 2001;153:114-22.
21. Hung HC, Huang MC, Lee JM, et al. Association between

- diet and esophageal cancer in Taiwan. *J Gastroenterol Hepatol* 2004;9:632-7.
22. De Stefani E, Deneo-Pellegrini H, Ronco AL, et al. Food groups and risk of squamous cell carcinoma of the oesophagus: a case-control study in Uruguay. *Br J Cancer* 2003;89:1209-14.
 23. Wang JM, Xu B, Rao JY, et al. Diet habits, alcohol drinking, tobacco smoking, green tea drinking, and the risk of esophageal squamous cell carcinoma in the Chinese population. *Eur J Gastroenterol Hepatol* 2007;19:171-6.
 24. De Stefani E, Ronco AL, Boffetta P, et al. Nutrient intake and risk of squamous cell carcinoma of the esophagus: a case-control study in Uruguay. *Nutr Cancer* 2006;56:149-57.
 25. Bollschweiler E, Wolfgarten E, Nowroth T, et al. Vitamin intake and risk of subtypes of esophageal cancer in Germany. *J Cancer Res Clin Oncol* 2002;128:575-80.
 26. Franceschi S, Bidoli E, Negri E, et al. Role of macronutrients, vitamins and minerals in the aetiology of squamous-cell carcinoma of the oesophagus. *Int J Cancer* 2000;86:626-31.
 27. Blot WJ, Li JY, Taylor PR, et al. Nutrition intervention trials in Linxian, China: supplementation with specific vitamin/mineral combinations, cancer incidence, and disease-specific mortality in the general population. *J Natl Cancer Inst* 1993;85:1483-92.
 28. Mark SD, Qiao YL, Dawsey SM, et al. Prospective study of serum selenium levels and incident esophageal and gastric cancers. *J Natl Cancer Inst* 2000;92:1753-63.
 29. Wei WQ, Abnet CC, Qiao YL, et al. Prospective study of serum selenium concentrations and esophageal and gastric cardia cancer, heart disease, stroke, and total death. *Am J Clin Nutr* 2004;79:80-5.
 30. Abnet CC, Lai B, Qiao Y-L, et al. Zinc concentration in esophageal biopsies measured by X-ray fluorescence and cancer risk. *J Natl Cancer Inst* 2005;97:301-6.
 31. Terry P, Lagergren J, Ye W, Nyrén O, Wolk A. Antioxidants and cancers of the esophagus and gastric cardia. *Int J Cancer* 2000;87:750-4.
 32. Kubo A, Corley DA. Meta-analysis of antioxidant intake and the risk of esophageal and gastric cardia adenocarcinoma. *Am J Gastroenterol* 2007;102:2323-30.
 33. González CA, Pera G, Agudo A, et al. Fruit and vegetable intake and the risk of stomach and oesophagus adenocarcinoma in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Int J Cancer* 2006;118:2559-66.
 34. González CA, Jakszyn P, Pera G, et al. Meat intake and risk of stomach and esophageal adenocarcinoma within the European Prospective Investigation Into Cancer and Nutrition (EPIC). *J Natl Cancer Inst* 2006;98:345-54.
 35. Corley DA, Kerlikowske K, Verma R, Buffler P. Protective association of aspirin/NSAIDs and esophageal cancer: a systematic review and meta-analysis. *Gastroenterology* 2003; 124: 47-56.
 36. Lindblad M, Lagergren J, García Rodríguez LA. Nonsteroidal anti-inflammatory drugs and risk of esophageal and gastric cancer. *Cancer Epidemiol Biomarkers Prev* 2005;14:444-50.
 37. Anderson LA, Johnston BT, Watson RG, et al. Nonsteroidal anti-inflammatory drugs and the esophageal inflammation-metaplasia-adenocarcinoma sequence. *Cancer Res* 2006; 66:4975-82.
 38. Fortuny J, Johnson CC, Bohlke K, et al. Use of anti-inflammatory drugs and lower esophageal sphincter-relaxing drugs and risk of esophageal and gastric cancers. *Clin Gastroenterol Hepatol* 2007;5:1154-9.
 39. Duan L, Wu AH, Sullivan-Halley J, Bernstein L. Nonsteroidal anti-inflammatory drugs and risk of esophageal and gastric adenocarcinomas in Los Angeles County. *Cancer Epidemiol Biomarkers Prev* 2008;17:126-34.
 40. Limburg PJ, Wei W, Ahnen DJ, et al. Randomized, placebo-controlled, esophageal squamous cell cancer chemoprevention trial of selenomethionine and celecoxib. *Gastroenterology* 2005;129:863-73.
 41. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. Human papillomaviruses. *IARC Monogr Eval Carcinog Risks Hum* 2007;90:1-636.
 42. Gillison ML, Shah KV. Chapter 9: Role of mucosal human papillomavirus in nongenital cancers. *J Natl Cancer Inst Monogr* 2003;31:57-65.
 43. Yao PF, Li GC, Li J, et al. Evidence of human papilloma virus infection and its epidemiology in esophageal squamous cell carcinoma. *World J Gastroenterol* 2006;12:1352-5.
 44. Koh JS, Lee SS, Baek HJ, Kim YI. No association of high-risk human papillomavirus with esophageal squamous cell carcinomas among Koreans, as determined by polymerase chain reaction. *Dis Esophagus* 2008;21:114-7.
 45. Talamini G, Capelli P, Zamboni G, et al. Alcohol, smoking and papillomavirus infection as risk factors for esophageal squamous-cell papilloma and esophageal squamous-cell carcinoma in Italy. *Int J Cancer* 2000;86:874-8.
 46. White RE, Mungatana C, Mutuma G, et al. Absence of human papillomavirus in esophageal carcinomas from southwestern Kenya. *Dis Esophagus* 2005;18:28-30.
 47. Gao GF, Roth MJ, Wei WQ, et al. No association between HPV infection and the neoplastic progression of esophageal squamous cell carcinoma: result from a cross-sectional study in a high-risk region of China. *Int J Cancer* 2006;119: 1354-9.
 48. International Agency for Research on Cancer. Human papillomaviruses. *IARC Monogr Eval Carcinog Risks Hum*. 2007;90:1-636.
 49. Nideobitek G. Epstein-Barr virus infection in the pathogenesis of nasopharyngeal carcinoma. *J Clin Pathol: Mol Pathol*, 53: 248-54.
 50. Wang LS, Chow KC, Wu YC, Li WY, Huang MH. Detection of Epstein-Barr virus in esophageal squamous cell carcinoma in Taiwan. *Am J Gastroenterol* 1999;94:2834-9.
 51. Awerkw S, Bollschweiler E, Metzger R, Schneider PM, Hölscher AH, Pfister H. Esophageal cancer in Germany is associated with Epstein-Barr-virus but not with papillomaviruses. *Med Microbiol Immunol* 2003;192:137-40.
 52. Sunpaweravong S, Mitarnun W, Puttawibul P. Absence of Epstein-Barr virus in esophageal squamous cell carcinoma.

- Dis Esophagus 2005;18:319-35.
53. Yanai H, Hirano A, Matsusaki K, et al. Epstein-Barr virus association is rare in esophageal squamous cell carcinoma. *Int J Gastrointest Cancer* 2003;33:165-70.
 54. Mizobuchi S, Sakamoto H, Tachimori Y, Kato H, Watanabe H, Terada M. Absence of human papillomavirus-16 and -18 DNA and Epstein-Barr virus DNA in esophageal squamous cell carcinoma. *Jpn J Clin Oncol* 1997;27:1-5.
 55. Hong T, Shimada Y, Kano M, et al. The Epstein-Barr virus is rarely associated with esophageal cancer. *Int J Mol Med* 2000;5:363-8.
 56. Chang F, Syrjänen S, Shen Q, et al. Evaluation of HPV, CMV, HSV and EBV in esophageal squamous cell carcinomas from a high-incidence area of China. *Anticancer Res.* 2000; 20: 3935-40.
 57. Wang J, Noffsinger A, Stemmermann G, Fenoglio-Preiser C. Esophageal squamous cell carcinomas arising in patients from a high-risk area of North China lack an association with Epstein-Barr virus. *Cancer Epidemiol Biomarkers Prev* 1999; 8:1111-4.
 58. Cho YJ, Chang MS, Park SH, Kim HS, Kim WH. In situ hybridization of Epstein-Barr virus in tumor cells and tumor-infiltrating lymphocytes of the gastrointestinal tract. *Hum Pathol* 2001;32:297-301.
 59. Peek RM, Crabtree JE. *Helicobacter* infection and gastric neoplasia. *J Pathol* 2006;208:233-48.
 60. Suerbaum S. and Michetti P. *Helicobacter pylori* infection. *N Engl. J Med* 2002;347:1175-86.
 61. Islami F, Kamangar F. *Helicobacter pylori* and esophageal cancer risk: a meta-analysis. *Cancer Prev Res (Phila Pa)* 2008;1:329-38.
 62. Rokkas T, Pistiolas D, Sechopoulos P, Robotis I, Margantinis G. Relationship between *Helicobacter pylori* infection and esophageal neoplasia: a meta-analysis. *Clin Gastroenterol Hepatol* 2007;5:1413-7.
 63. Zhuo X, Zhang Y, Wang Y, Zhuo W, Zhu Y, Zhang X. *Helicobacter pylori* infection and oesophageal cancer risk: association studies via evidence-based meta-analyses. *Clin Oncol (R Coll Radiol)* 2008;20:757-62.
 64. Chow WH, Blaser MJ, Blot WJ, et al. An inverse relation between cagA+ strains of *Helicobacter pylori* infection and risk of esophageal and gastric cardia adenocarcinoma. *Cancer Res* 1998;58:588-90.
 65. Fong LY, Lau KM, Huebner K, Magee PN. Induction of esophageal tumors in zinc-deficient rats by single low doses of N-nitrosomethylbenzylamine (NMBA): analysis of cell proliferation, and mutations in H-ras and p53 genes. *Carcinogenesis* 1997;18:1477-84.
 66. Lijinsky W, Singer GM, Kovatch RM. Similar carcinogenic effects in rats of 1-ethyl-1-nitroso-3-hydroxyethylurea and 1-hydroxyethyl-1-nitroso-3-ethylurea. *Carcinogenesis* 1985;6: 641-3.
 67. Ivankovic S, Seibel J, Komitowski D, Spiegelhalter B, Preussmann R, Siddiqi M. Caffeine-derived N-nitroso compounds. V. Carcinogenicity of mononitrosocaffeidine and dinitrosocaffeidine in bd-ix rats. *Carcinogenesis* 1998; 19:933-7.
 68. Preussmann R, Habs M, Habs H, Stummeyer D. Fluoro-substituted N-nitrosamines. 6. carcinogenicity of N-nitroso-(2,2,2-trifluoroethyl)-ethylamine in rats. *Carcinogenesis* 1983; 4:755-7.
 69. Bartsch H, Frank N, Frei E, Marks F, Schroeder CH, Lin JK. Taiwanese-German workshop on tumour prevention. *Eur J Cancer Prev* 1996;5:83-8.
 70. Tricker AR. N-nitroso compounds and man: sources of exposure, endogenous formation and occurrence in body fluids. *Eur J Cancer Prev* 1997;6:226-68.
 71. Abnet CC, Qiao YL, Mark SD, Dong ZW, Taylor PR, Dawsey SM. Prospective study of tooth loss and incident esophageal and gastric cancers in China. *Cancer Causes Control* 2001; 12:847-54.
 72. Abnet CC, Kamangar F, Dawsey SM, et al. Tooth loss is associated with increased risk of gastric non-cardia adenocarcinoma in a cohort of Finnish smokers. *Scand J Gastroenterol* 2005;40:681-7.
 73. Jakszyn P, Gonzalez CA. Nitrosamine and related food intake and gastric and oesophageal cancer risk: a systematic review of the epidemiological evidence. *World J Gastroenterol* 2006;12:4296-303.
 74. Danaei G, Vander Hoorn S, Lopez AD, Murray CJ, Ezzati M. Comparative Risk Assessment collaborating group (Cancers). Causes of cancer in the world: comparative risk assessment of nine behavioural and environmental risk factors. *Lancet* 2005;366:1784-93.
 75. Kazerouni N, Sinha R, Hsu CH, Greenberg A, Rothman N. Analysis of 200 food items for benzo(a)pyrene and estimation of its intake in an epidemiologic study. *Food Chem Toxicol* 2001;39:423-36.
 76. Chang KF, Fang GC, Chen JC, Wu YS. Atmospheric polycyclic aromatic hydrocarbons (PAHs) in Asia: a review from 1999 to 2004. *Environ Pollut* 2006;142:388-96.
 77. Boffetta P, Jourenkova N, Gustavsson P. Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes Control* 1997;8: 444-72.
 78. Roth MJ, Guo-Qing W, Lewin KJ, et al. Histopathologic changes seen in esophagectomy specimens from the high-risk region of Linxian, China: potential clues to an etiologic exposure? *Hum Pathol* 1998;29:1294-8.
 79. Roth MJ, Strickland KL, Wang GQ, Rothman N, Greenberg A, Dawsey SM. High levels of carcinogenic polycyclic aromatic hydrocarbons present within food from Linxian, China may contribute to that region's high incidence of oesophageal cancer. *Eur J Cancer* 1998;34:757-8.
 80. Roth MJ, Qiao YL, Rothman N, et al. High urine 1-hydroxypyrene glucuronide concentrations in Linxian, China, an area of high risk for squamous oesophageal cancer. *Biomarkers* 2001;6:381-6.
 81. Strickland P, Kang D, Sithisarankul P. Polycyclic aromatic hydrocarbon metabolites in urine as biomarkers of exposure and effect. *Environ Health Perspect* 1996;104 (Suppl 5): 927-32.

82. Yokoyama A, Omori T. Genetic polymorphisms of alcohol and aldehyde dehydrogenases and risk for esophageal and head and neck cancers. *Alcohol* 2005;35:175-85.
83. International Agency for Research on Cancer. Acetaldehyde. IARC Monogr Eval Carcinog Risks Hum 1999;71Pt2:319-35.
84. Lagergren J, Bergström R, Lindgren A, Nyrén O. Symptomatic gastroesophageal reflux as a risk factor for esophageal adenocarcinoma. *N Engl J Med* 1999;340:825-31.
85. Wu AH, Tseng CC, Bernstein L. Hiatal hernia, reflux symptoms, body size, and risk of esophageal and gastric adenocarcinoma. *Cancer* 2003;98:940-8.
86. Farrow DC, Vaughan TL, Sweeney C, et al. Gastroesophageal reflux disease, use of H2 receptor antagonists, and risk of esophageal and gastric cancer. *Cancer Causes Control* 2000;11:231-8.
87. Shaheen NJ, Crosby MA, Bozymski EM, Sandler RS. Is there publication bias in the reporting of cancer risk in Barrett's esophagus? *Gastroenterology* 2000;119:333-8.
88. Jankowski JA, Provenzale D, Moayyedi P. Esophageal adenocarcinoma arising from Barrett's metaplasia has regional variations in the west. *Gastroenterology* 2002;122:588-90.
89. Lieberman DA, Oehlke M, Helfand M. Risk factors for Barrett's esophagus in community-based practice. GORGE consortium. Gastroenterology Outcomes Research Group in Endoscopy. *Am J Gastroenterol* 1997;92:1293-7.
90. Ryan AM, Rowley SP, Fitzgerald AP, Ravi N, Reynolds JV. Adenocarcinoma of the oesophagus and gastric cardia: male preponderance in association with obesity. *Eur J Cancer* 2006;42:1151-8.
91. Merry AH, Schouten LJ, Goldbohm RA, van den Brandt PA. Body mass index, height and risk of adenocarcinoma of the oesophagus and gastric cardia: a prospective cohort study. *Gut* 2007;56:1503-11.
92. Kubo A, Corley DA. Body mass index and adenocarcinomas of the esophagus or gastric cardia: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15:872-8.
93. Hampel H, Abraham NS, El-Serag HB. Meta-analysis: obesity and the risk for gastroesophageal reflux disease and its complications. *Ann Intern Med* 2005;143:199-211.
94. Corley DA, Levin TR, Habel LA, Buffler PA. Barrett's esophagus and medications that relax the lower esophageal sphincter. *Am J Gastroenterol* 2006;101:937-44.
95. Lindblad M, Rodríguez LA, Lagergren J. Body mass, tobacco and alcohol and risk of esophageal, gastric cardia, and gastric non-cardia adenocarcinoma among men and women in a nested case-control study. *Cancer Causes Control* 2005;16:285-94.
96. Jansson C, Johansson AL, Nyrén O, Lagergren J. Socio-economic factors and risk of esophageal adenocarcinoma: a nationwide Swedish case-control study. *Cancer Epidemiol Biomarkers Prev* 2005;14:1754-61.
97. Brown LM, Devesa SS. Epidemiologic trends in esophageal and gastric cancer in the United States. *Surg Oncol Clin N Am* 2002;11:235-56.