Review Article • บทฟื้นฟูวิชาการ

Electrolytes: Unraveling their Critical Importance in Anesthesia Practice

Krongthip Sripunjan, Napichayakarn Phanthananphokhin, Wilaiporn Supan, Phongthara Vichitvejpaisal

Department of Anesthesiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

Abstract

This article explores the critical significance of electrolytes in anesthesia practice and their impact on patient outcomes. It provides a comprehensive overview of electrolytes and their role in the human body, emphasizing their essential functions in maintaining physiological processes. It highlights the implications of electrolyte imbalances during the perioperative period and their potential effects on cardiac function, neuromuscular function, wound healing, and organ function. Strategies for monitoring, managing, and optimizing electrolyte balance in anesthesia practice are discussed. The article also presents the critical importance of maintaining electrolyte balance and the need for timely interventions during anesthesia practice. Additionally, future directions in electrolyte monitoring, emerging research areas, and the critical role of electrolytes in anesthesia practice and patient outcomes are explored.

Keywords: anesthesia practice, electrolytes, homeostasis, intravenous fluid

Key points

- Electrolytes are essential minerals that carry an electric charge and play a vital role in maintaining physiological processes in the human body. Maintaining proper electrolyte balance is crucial in anesthesia practice to prevent adverse effects on patient outcomes, including cardiac arrhythmias, neuromuscular dysfunction, and impaired organ function.
- Preoperative assessment and optimization of electrolyte levels are necessary to identify existing imbalances and correct them before surgery. Intraoperative electrolyte management involves appropriate fluid selection, electrolyte supplementation, and continuous monitoring to maintain optimal electrolyte levels. Postoperative electrolyte monitoring is crucial to detect and address any imbalances that may impact patient recovery and outcomes.
- Electrolytes play a critical role in anesthesia practice, and prioritizing electrolyte balance is essential for patient safety and positive outcomes. Case studies and clinical examples illustrate the importance of early recognition and intervention in managing electrolyte disturbances during anesthesia practice.
- Future directions in electrolyte monitoring include advancements in point-of-care testing, wearable sensors, and the integration of artificial intelligence and data analytics. Emerging research areas focus on individualized electrolyte management, the impact of electrolyte balance on anesthetic pharmacology, and optimizing fluid resuscitation strategies.

Correspondence to: Phongthara Vichitvejpaisal, E-mail: phongthara@gmail.com

Received August 3, 2023, Revised December 13, 2023, Accepted December 13, 2023

Introduction

Electrolytes are essential minerals that carry an electric charge when dissolved in bodily fluids. They play a vital role in maintaining various physiological processes in the human body. Common electrolytes include sodium, potassium, calcium, magnesium, chloride, and bicarbonate.[1] These electrolytes are involved in the regulation of fluid balance, nerve function, muscle contractions, pH balance, and other crucial cellular activities. In the field of anesthesia, maintaining proper electrolyte balance is of utmost importance. Anesthesia procedures can have significant effects on electrolyte levels due to various factors, such as the administration of intravenous fluids, changes in fluid balance, altered metabolism, and physiological stress induced by surgical procedures. Electrolyte imbalances can lead to adverse effects on patient outcomes, including cardiac arrhythmias, neuromuscular dysfunction, impaired organ function, and delayed recovery.

The purpose of this article is to delve into the critical importance of electrolytes in anesthesia practice and highlight their significance in patient care. By exploring the relationship between electrolyte balance and anesthesia, the article aims to shed light on the potential implications of electrolyte disturbances during perioperative periods. It will emphasize the role of anesthesiologists and healthcare providers in monitoring, managing, and optimizing electrolyte levels to enhance patient safety, surgical outcomes, and recovery. Furthermore, the article will provide an analysis of various electrolyte disorders commonly encountered in anesthesia practice, such as hypernatremia, hyponatremia, hypokalemia, and hyperkalemia. It will discuss their causes, effects on anesthesia, and appropriate management strategies.

Understanding electrolytes

Electrolytes are minerals that carry an electric charge when dissolved in water or bodily fluids. They are categorized based on their positive or negative charge: Cations are positively charged electrolytes, e.g., sodium (Na^+), potassium (K^+), calcium (Ca^{++}), and magnesium (Mg^{++}). Anions are negatively charged electrolytes, e.g., chloride (Cl^-) and bicarbonate (HCO_3^-). These electrolytes are present in various body fluids, including blood, plasma, intracellular fluid, and interstitial fluid.[2]

Electrolyte balance and homeostasis

Maintaining electrolyte balance is crucial for optimal cellular function and overall health. The body tightly regulates electrolyte levels through complex mechanisms to ensure homeostasis. Electrolytes are obtained through dietary intake. The digestive system absorbs them into the bloodstream. They are distributed throughout the body via the circulatory system. Specialized transport proteins facilitate their movement across cell membranes. Excess electrolytes are eliminated through urine, feces, sweat, and respiration.[3] The kidneys play a vital role in regulating electrolyte excretion. Hormones, such as aldosterone, antidiuretic hormone (ADH), and parathyroid hormone (PTH), help regulate electrolyte levels by influencing renal reabsorption and excretion.

Electrolyte imbalances and their potential impact

Disruptions in electrolyte balance can have significant implications for patient health. Electrolyte imbalances can impact various bodily systems, including the cardiovascular, nervous, muscular, and renal systems. They can cause cardiac arrhythmias, muscle weakness, altered mental status, impaired organ function, and electrolyte-induced organ damage. Electrolyte imbalances can arise due to various factors, including medical conditions, medications, diet, and fluid imbalances.

Hyponatremia can lead to symptoms such as fatigue, confusion, muscle cramps, seizures, and in severe cases, brain swelling.[4]

Hypernatremia can cause thirst, dry mucous membranes, restlessness, altered mental status, and, in severe cases, neurological complications.[5]

Hypokalemia can result in muscle weakness, irregular heart rhythms, fatigue, constipation, and impaired kidney function.[6]

Hyperkalemia can lead to cardiac arrhythmias, muscle weakness, numbness, tingling, and potentially life-threatening conditions.[7]

Hypocalcemia can cause muscle spasms, numbness and tingling, impaired blood clotting, and weakened bones.[8]

Hypercalcemia can result in excessive thirst, kidney stones, bone pain, confusion, and cardiac abnormalities.[8]

Electrolytes and anesthesia

Anesthetic procedures can have a significant impact on electrolyte levels in the body. Since intravenous fluids are commonly administered during anesthesia to maintain hydration and compensate for fluid losses. These fluids can contain electrolytes that may affect the overall electrolyte balance. In addition, surgical procedures, especially those involving blood loss or fluid shifts, can cause changes in electrolyte concentrations. For example, extensive fluid losses during surgery can lead to electrolyte imbalances.

Anesthesia can also affect metabolic processes, including glucose metabolism and hormonal regulation, which can influence electrolyte levels. For instance, the stress response induced by anesthesia can trigger hormonal changes that impact electrolyte balance.[9] Moreover, anesthetic medications and adjuncts, such as diuretics, can also impact electrolyte levels. Certain medications may affect renal function and electrolyte excretion, potentially leading to imbalances.

Preoperative assessment

Before surgery, a thorough preoperative assessment is conducted, reviewing the patient's medical history, medications, and laboratory results to identify any existing electrolyte imbalances. To enhance early detection of electrolyte disturbances, patient education is prioritized, stressing the significance of maintaining proper electrolyte balance[10]., adhering to dietary considerations, and promptly reporting any symptoms or changes indicating electrolyte imbalances. Active patient involvement in their care further supports this process. Adequate hydration is ensured, and any fluid imbalances, like dehydration or overhydration, are promptly corrected to promote optimal electrolyte

balance. Anesthesiologists also evaluate the patient's baseline electrolyte levels through blood tests before surgery, which aids in identifying any existing electrolyte imbalances or conditions that may make the patient susceptible to electrolyte disturbances.

If electrolyte imbalances are found, appropriate measures are taken to correct them before the surgical procedure, which may involve dietary adjustments, oral or intravenous electrolyte supplementation[11], or specific medical interventions. Close collaboration with the surgical team and healthcare professionals is vital throughout the perioperative period. By addressing existing electrolyte imbalances, the surgical team can help restore electrolyte balance, reducing the risk of perioperative complications and improving patient outcomes.

Intraoperative interventions

One crucial aspect of patient management during procedures involves the careful selection of intravenous fluids tailored to their specific needs. Among the options available are crystalloids and colloids[12], each with their unique electrolyte compositions, which can be utilized based on the patient's requirements. During the procedure, vigilant monitoring of electrolyte levels and fluid balance is of utmost importance. This monitoring is accomplished through laboratory tests or point-of-care testing, allowing healthcare providers to swiftly identify and address any electrolyte imbalances that may arise. By administering appropriate electrolyte supplementation and adjusting fluid rates and composition as necessary, optimal electrolyte levels can be maintained, taking into account factors such as ongoing losses, surgical duration, and electrolyte imbalances detected during monitoring.

As the surgical process unfolds, changes may occur in factors like blood loss or fluid shifts. It is essential for the fluid management plan to remain dynamic, adapting to these alterations and ensuring the patient's well-being is continually safeguarded. The monitoring of electrolyte levels during anesthesia procedures is typically carried out through blood tests. These tests, which can be performed periodically or through intraoperative blood gas analysis, furnish real-time information on electrolyte concentrations. However, in certain situations, particularly when time is of the essence, point-of-care testing devices prove to be invaluable.

Postoperative monitoring

Obtain postoperative blood samples to assess electrolyte levels and identify any imbalances or changes compared to baseline measurements. If postoperative electrolyte imbalances are detected, provide appropriate electrolyte supplementation to restore and maintain optimal levels. This may involve oral or intravenous electrolyte administration, depending on the severity of the imbalance.[13] Monitor patients closely for signs and symptoms of electrolyte disturbances during the recovery phase. Educate patients about the importance of electrolyte balance, signs of imbalances, and the need to report any abnormalities to healthcare providers. Maintain open communication with the surgical team, nursing staff, and other healthcare providers involved in postoperative care to ensure coordinated efforts in managing electrolyte balance and addressing any electrolyte-related

concerns promptly.

Specific anesthetic techniques

Certain anesthetic techniques and interventions may have specific implications for electrolyte balance. Different fluid replacement strategies, such as crystalloids and colloids, may have varying effects on electrolyte concentrations. Anesthesiologists must consider the composition of fluids administered to ensure electrolyte balance is maintained appropriately. Significant blood loss during surgery can also lead to electrolyte imbalances[14], particularly if blood transfusions are required. Packed red blood cell transfusions may affect electrolyte concentrations, necessitating monitoring and appropriate supplementation.

Induced hypothermia, used in certain surgical procedures, can impact electrolyte homeostasis. Hypothermia may alter cellular metabolism, enzyme activity, and ion channel function, potentially affecting electrolyte levels. Additionally, diuretic medications, sometimes used during anesthesia, can promote diuresis and alter electrolyte excretion. Careful monitoring is necessary to prevent excessive electrolyte losses.[15]

Electrolytes and surgical outcomes

Electrolyte imbalances can significantly impact perioperative complications and patient outcomes. Hypokalemia or hyperkalemia, can lead to cardiac arrhythmias, including ventricular tachycardia or fibrillation. These abnormalities can result in hemodynamic instability, myocardial ischemia, or even cardiac arrest. Imbalances in electrolytes like sodium, potassium, and magnesium can cause neuromuscular dysfunction[16], leading to muscle weakness, impaired respiratory function, and postoperative complications such as respiratory failure.

Electrolyte imbalances, particularly hypocalcemia, can hinder the process of wound healing.[17] Calcium plays a critical role in cell migration, collagen synthesis, and tissue repair. Insufficient calcium levels can delay wound healing and increase the risk of wound complications. Moreover, electrolyte imbalances may affect various organ systems. For instance, electrolyte disturbances can impair renal function, leading to acute kidney injury or delayed recovery of renal function postoperatively.[18]

Postoperative recovery

Optimal electrolyte balance is crucial for postoperative recovery and patient well-being. Maintaining appropriate electrolyte levels contributes to overall fluid balance. Adequate fluid balance is essential for organ perfusion, tissue oxygenation, and efficient metabolic processes, all of which are critical for postoperative recovery. Electrolytes are involved in various metabolic pathways, including energy production, enzyme activity, and acid-base balance.[19] Optimal electrolyte levels support metabolic processes necessary for wound healing, tissue repair, and cellular regeneration.

Notably, electrolyte imbalances can result in muscle weakness, which can impede early mobilization and delay the recovery process. Restoring electrolyte balance helps promote muscle strength and mobility. Furthermore,

abnormalities in sodium and magnesium, can affect neurological function, leading to cognitive impairment, confusion, and altered mental status.20 Maintaining electrolyte balance is crucial for postoperative neurological recovery.

Electrolyte disorders and anesthetic considerations

Hypernatremia

Hypernatremia can be caused by factors such as excessive sodium intake, inadequate water intake, diabetes insipidus, or certain medical conditions.[21] Patients at risk include those with impaired thirst mechanisms, limited access to water, or altered fluid balance regulation. Patients with severe hypernatremia may present with dehydration, altered sensorium, or neurological manifestations.[5] Treatment involves addressing the underlying cause and correcting the fluid imbalance. Oral or intravenous administration of isotonic or hypotonic fluids may be necessary to restore water balance and lower sodium levels gradually.[14]

Hyponatremia

Hyponatremia can result from conditions such as excessive fluid intake, fluid overload, certain medications, or underlying medical conditions. It can lead to neurological symptoms, including confusion, seizures, and cerebral edema. During anesthesia, hyponatremia may increase the risk of cerebral edema, alter drug metabolism, and affect neuromuscular function. Monitoring of electrolyte levels, fluid status, and hemodynamic parameters is crucial. Close neurological monitoring and appropriate management of cerebral edema are essential to prevent complications.[4] Treatment aims to correct the underlying cause of hyponatremia and normalize sodium levels while carefully managing fluid balance. Interventions may include fluid restriction, diuretic therapy, or hypertonic saline infusion based on the severity and clinical presentation.

Hypokalemia and hyperkalemia

Hypokalemia can arise from conditions such as gastrointestinal losses, diuretic use, renal disorders, or certain medications. Instead, hyperkalemia can result from factors such as renal dysfunction, acidosis, tissue injury, or certain medications. However, both can lead to cardiac arrhythmias, muscle weakness, and impaired neuromuscular function. Close monitoring of cardiac function, including ECG monitoring, and potassium levels is crucial.[6,22] For hypokalemia, potassium supplementation may be necessary before surgery to restore normal levels and minimize the risk of arrhythmias or neuromuscular complications. On the other hand, treatment approaches for hyperkalemia may include calcium administration to stabilize cardiac membranes, insulin and glucose therapy to shift potassium intracellularly, and measures to enhance potassium elimination.[6,7]

Future directions

The development of advanced point-of-care testing devices allows for rapid, accurate, and convenient electrolyte measurements at the patient's bedside. These devices can provide real-time results, enabling prompt interventions

and adjustments in electrolyte management. Emerging technologies such as wearable sensors have the potential to revolutionize electrolyte monitoring. These non-invasive devices can continuously monitor electrolyte levels and provide real-time data, enhancing patient safety and allowing for early detection of imbalances. Integration of artificial intelligence (AI) and data analytics into electrolyte management systems can aid in predicting and preventing electrolyte imbalances.[23]

Further investigation is needed to understand how electrolyte imbalances influence the pharmacokinetics and pharmacodynamics of anesthetic drugs. This knowledge can guide dosage adjustments and optimize drug efficacy and safety. Additionally, research is exploring the impact of different fluid resuscitation strategies on electrolyte balance, including the use of balanced crystalloids, colloids, and individualized fluid regimens.[24] The goal is to optimize fluid management and minimize electrolyte disturbances.

Conclusion

Electrolytes play a critical role in anesthesia practice, and maintaining optimal electrolyte balance is essential for patient safety and positive outcomes. Electrolyte imbalances can have significant implications, including cardiac arrhythmias, neuromuscular dysfunction, impaired wound healing, and organ dysfunction. Anesthesiologists must prioritize electrolyte monitoring, recognize imbalances, and implement appropriate interventions to maintain electrolyte homeostasis. The future of electrolyte management in anesthesia practice holds promising advancements, including improved monitoring technologies, personalized approaches, and integration of AI and data analytics. Ongoing research is expanding our knowledge of electrolyte-related anesthesia considerations, optimizing pharmacological approaches, and refining fluid resuscitation strategies.

References

1. Shrimanker I, Bhattarai S. Electrolytes. StatePearls. 2023 [update 2023 Apr 23]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK541123/.
2. Rassam SS, Counsell DJ. Perioperative electrolyte and fluid balance. Cont Educ Anaesth Crit Care Pain. 2005;5:157-60.
3. Morris AL, Mohiuddin SS. Biochemistry, Nutrients. StatPreals. 2023 [update 2023 May 1; cited 2023 Jul 28]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK554545/
4. Leung AA, McAlister FA, Rogers SO, Pazo V, Wright A, Bates DW. Preoperative hyponatremia and perioperative complications. Arch Intern Med. 2012;172:1474-81.
5. Qian Q. Hypernatremia. Clin J Am Soc Nephrol. 2019;14:432-4.
6. Castro D, Sharma S. Hypokalemia. StatPearls. 2023 [cited 2023 Jul 28]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK482465/.
7. Simon LV, Hashmi MF, Farrell MW. Hyperkalemia. StatPearls. 2023 [update 2023 Feb 19; cited 2023 Jul 28]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK4702843/.
8. Diringer M. Neurologic manifestations of major electrolyte abnormalities. Handbook of clinical neurology. 2017 [cited 2023 Jul 28]. p705-13. Available from: https://pubmed.ncbi.nlm.nih.gov/28190443/.
9. Hahn RG. Fluid and electrolyte physiology in anaesthetic practice. In: Hardman JG, Hardman JG, Hopkins PM, Struys MMRF, editors. Oxford Textbook of Anaesthesia: Oxford University Press.

2017 [cited 2023 Jul 28]. p49-58. Available from: https://academic.oup.com/book/30062/chapter-abstract/256303597?redirectedFrom=fulltext&login=true.
10. Mauricio Del Rio J, Nicoara A, Swaminathan M. Neuroendocrine stress response: implication for cardiac surgery-associated acute kidney injury. Rom J Anesth Intensive Care. 2017;24:57-63.
11. Wagner J. The positives and negatives of preoperative electrolyte disturbances. South African J Anaesth Analg. 2021;27:126-30.
12. Marx G, Meybohm P, Schuerholz T, et al. Impact of a new balanced gelatine on electrolytes and pH in the perioperative care. PLoS One. 2019;14:1-15.
13. National Clinical Guideline Center. Intravascular fluid therapy: intravenous fluid therapy in adults in hospital. London: Royal College of Physicians; 2013 [cited 2023 Jul 28]. Available from: https://pubmed.ncbi.nlm.nih.gov/25340240/.
14. Muhsin SA, Mount DB. Diagnosis and treatment of hypernatremia. Best Pract Res Clin Endocrinol Metab. 2016;30:189-203.
15. Arampatzis S, Funk GC, Leichtle AB, et al. Impact of diuretic therapy-associated electrolyte disorders present on admission to the emergency department: a cross-sectional analysis. BMC Med. 2013;11:1-6.
16. Wenwen X, Jing Y, Yingchao S, Qinglu W. The effect of magnesium deficiency on neurological disorders: a narrative review article. Iran J Public Health. 2019;48:379-87.
17. Subramaniam T, Fauzi MB, Lokanathan Y, Law JX. The role of calcium in wound healing. Int J Mol Sci. 2021;22:6486.
18. Goyal A, Daneshpajouhnejad P, Hashmi MF, Bashir K. Acute kidney injury. StatPearls. 2023 [cited 2023 Jul 28]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK441896/
19. Simpson R, Quayle J, Stylianides N, Carlson G, Soop M. Intravenous fluid and electrolyte administration in elective gastrointestinal surgery: mechanisms of excessive therapy. Ann R Coll Surg Engl. 2017;99:497-503.
20. Wagner J. The positives and negatives of preoperative electrolyte disturbances. South African J Anaesth Analg. 2021;27:126-30.
21. Bennet D, Khorsandian Y, Pelusi J, Mirabella A, Pirrotte P, Zenhausern F. Molecular and physical technologies for monitoring fluid and electrolyte imbalance: a focus on cancer population. Clin Transl Med. 2021;11:1-22.
22. Freshwater DT. Sodium, potassium and the anaesthetist. World Anaesthesia Tutorial of the week. 2005 [cited 2023 Jul 28]. Available from: https://resources.wfsahq.org/atotw/sodium-potassium-and-the-anaesthetist/
23. Kwon JM, Jung MS, Kim KH, et al. Artificial intelligence for detecting electrolyte imbalance using electrocardiography. Ann Noninvasive Electrocardiol. 2021;26: e12839.
24. McClelland M. IV therapies for patients with fluid and electrolyte imbalances. Medsurg Nurs. 2014;23:S4-8.