

Advancements in Ambulatory Anesthesia Pain Management

Nuthakun Posri¹, Varathip Thongdech², Peeraya Buakhao³, Phongthara Vichitvejpaisal⁴

¹ Division of Perioperative Nursing, Golden Jubilee Medical Center,
Faculty of Medicine Siriraj Hospital, Mahidol University, Nakhon Pathom 73170, Thailand

² Division of Nursing Specialty, Golden Jubilee Medical Center,
Faculty of Medicine Siriraj Hospital, Mahidol University, Nakhon Pathom 73170, Thailand

³ Division of Emergency Medicine Nursing, Golden Jubilee Medical Center,
Faculty of Medicine Siriraj Hospital, Mahidol University, Nakhon Pathom 73170, Thailand

⁴ Department of Anesthesiology, Faculty of Medicine Siriraj Hospital, Mahidol University,
Bangkok 10700, Thailand

Abstract

This article discusses the advancements in ambulatory anesthesia pain management and their implications for modern healthcare. The importance of effective pain management in ambulatory settings is emphasized, along with the challenges of traditional pain management techniques. The article then focuses on multi-modal analgesia in ambulatory anesthesia, explaining its rationale and detailing the various drugs and techniques used. Furthermore, the article discusses the role of technology in ambulatory anesthesia pain management. It focuses on advancements in pain assessment tools, drug delivery systems, and the use of virtual reality as a distraction technique. The

integration of patient-centered approaches in pain management is also explored emphasizing individualized pain management plans and patient engagement. The article also addresses the management of chronic pain in ambulatory anesthesia patients, emphasizing the need to identify and develop specialized pain management strategies for this population. It also discusses strategies for addressing safety and complications in ambulatory anesthesia pain management. The future directions of ambulatory anesthesia pain management are explored, with a focus on potential areas for further research and advancements in the field.

Keywords: ambulatory anesthesia, multimodal analgesia, pain management, patient-centered care

Key points:

- Ambulatory anesthesia plays a significant role in modern healthcare by providing efficient and cost-effective care.
- Multimodal analgesia in ambulatory anesthesia offers enhanced pain relief, reduced opioid consumption, and faster recovery and discharge.
- Management of chronic pain in ambulatory settings requires specialized strategies and collaboration between anesthesia teams and pain specialists.

Introduction

Ambulatory anesthesia, also known as outpatient or day surgery anesthesia, is a specialized approach that allows patients to undergo surgical or medical procedures and return home on the same day. With the advent of ambulatory anesthesia, a wide range of surgeries and medical interventions can now be performed in dedicated ambulatory surgery centers or outpatient facilities. The main advantage of ambulatory anesthesia is that it allows patients to recover in the comfort of their own homes, promoting faster recuperation and a quicker return to daily activities. Additionally, it has evolved from the need to reduce hospital stays, improve patient comfort, streamline healthcare services, and enhance cost-effectiveness. As patients return home shortly after their procedures, they need to be adequately relieved of pain to ensure a smooth recovery process. Proper pain management enables patients to be more functional, perform necessary activities,

and adhere to post-operative instructions effectively. It is crucial to provide individualized pain management plans tailored to each patient's needs to achieve optimal results and overall patient satisfaction.

Evolution of ambulatory anesthesia

The evolution of ambulatory anesthesia has been a significant milestone in the history of medical care, revolutionizing pain management practices for outpatient surgeries and procedures. One of the pioneering advancements was the introduction of local anesthesia, which enabled surgeons to perform minor procedures without the need for general anesthesia. Local anesthetics, such as lidocaine, were administered at specific sites, providing targeted pain relief and allowing patients to be awake during the procedure. In the mid-20th century, with the advent of intravenous anesthetics and advancements in monitoring technologies, the concept of monitored anesthesia care (MAC) gained prominence.^{1,2} MAC allowed for a lighter level of sedation while ensuring patient comfort and safety. This approach was particularly suitable for certain outpatient procedures, as it allowed patients to recover more rapidly and return home sooner.² In the 1980s and 1990s, anesthesia providers began to develop tailored techniques for ambulatory settings, recognizing the need for effective pain management that allowed patients to resume normal activities promptly.³

Enhanced recovery after surgery (ERAS)

ERAS is a comprehensive, evidence-based

approach to perioperative care designed to optimize patient outcomes, expedite recovery, and reduce hospital stays.⁴ Originally developed for major surgical procedures, ERAS encompasses various strategies, including preoperative optimization, patient education, minimization of surgical stress, effective pain management, early mobilization, and a well-coordinated, multi-disciplinary approach to care.⁵ Additionally, in ambulatory anesthesia, ERAS has been adapted to streamline the perioperative process, ensuring patients experience minimal disruption to their daily lives while recovering safely and comfortably at home. One of the key emphases of ERAS is the utilization of minimally invasive surgical approaches whenever feasible. These techniques are associated with reduced tissue damage, less post-operative pain, and a lower likelihood of complications. Consequently, patients benefit from faster recovery, shorter hospital stays, and increased satisfaction with the overall surgical experience.

A crucial aspect of ERAS is patient education, which plays a vital role in the successful implementation of the approach. Patients are provided with comprehensive information about the upcoming procedure, expected pain levels, and their role in managing pain during the recovery period. This education includes guidance on proper nutrition and hydration, the use of multimodal analgesia for effective pain control, and the potential benefits of non-pharmacological pain relief techniques. Encouraging early ambulation and physical activity post-surgery is also emphasized as part of the education process. ERAS extends its impact beyond the hospital setting, as it includes

post-discharge follow-up and support. Healthcare providers closely monitor patient progress, address any concerns, and offer guidance to optimize recovery outcomes. This ongoing support ensures that patients continue to manage pain effectively at home, ultimately contributing to successful long-term recovery. As a result of its comprehensive approach and focus on patient education and engagement, ERAS proves to be cost-effective for both patients and healthcare facilities. By promoting better outcomes and faster recovery times, ERAS improves patient satisfaction and overall healthcare efficiency.⁵

Individualized pain management

Individualized pain management recognizes the intricate interplay of emotional and physical discomfort influenced by diverse factors. The International Association for the Study of Pain defines pain as a distressing encounter involving sensory, emotional, cognitive, and social dimensions, often tied to tissue damage.⁶ Acute postoperative pain, originating from surgical tissue trauma, varies based on surgical type and peaks within the initial 3 days.⁷ The body's responses, orchestrated by the sympathetic and parasympathetic nervous systems, can trigger symptoms like elevated heart rate, increased blood pressure, or bronchospasm. These reactions might contribute to chronic pain and even metabolic alterations such as insulin resistance. Effective pain management, particularly in ambulatory surgery, holds pivotal importance for both the surgical procedure itself and immediate postoperative care. Such management empowers patients to adeptly manage pain in a home setting. Patient-centered approaches

to ambulatory anesthesia pain management acknowledge the idiosyncratic nature of each individual's pain experience and their responsiveness to pain management techniques. Thus, crafting personalized pain management plans tailored to each patient's unique requirements and attributes is paramount.

Diverse factors-ranging from pain tolerance and medical history to allergies and prior encounters with anesthesia and pain medications-demand meticulous consideration while devising pain management strategies. Customizing pain relief plans to suit the individual enables healthcare providers to optimize pain control, mitigate the risk of adverse effects, and amplify overall patient contentment. A fundamental facet of these patient-centered strategies involves collaborative decision-making between healthcare providers and patients. Engaging patients in the treatment planning process and factoring in their preferences and values empowers them to actively partake in their care, nurturing a sense of authority and independence. When patients perceive their contributions as valuable, adherence to the pain management plan improves, leading to superior pain control and enhanced treatment outcomes overall.

Multimodal analgesia

Multimodal analgesia, also known as balanced analgesia or combination analgesia, is an approach to pain management that involves using multiple drugs and techniques with different mechanisms of action to provide effective pain relief while minimizing the reliance

on opioids.⁸ The rationale behind multimodal analgesia lies in the concept of synergistic pain control, where each component of the analgesic regimen targets different pain pathways, producing a cumulative effect that enhances overall pain relief.

Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen and ketorolac, inhibit prostaglandin synthesis, which is involved in the inflammatory response and pain perception. By reducing inflammation, these drugs provide effective pain relief and are particularly useful for managing mild to moderate pain.⁹ COX-2 inhibitors, a specific type of NSAID, have the advantage of providing pain relief without interfering with blood clotting (unlike traditional NSAIDs that also inhibit COX-1). This characteristic makes them suitable for patients at risk of bleeding complications.¹⁰ Acetaminophen is an analgesic and antipyretic agent that can be used in combination with NSAIDs.¹¹ It acts centrally to modulate pain perception and fever.

Local anesthetics, such as lidocaine and bupivacaine, are administered to specific nerve or tissue sites to provide targeted pain relief. These can be used as part of regional anesthesia techniques, such as nerve blocks or epidurals, to block pain signals from reaching the brain. Drugs that antagonize the N-methyl-D-aspartate receptors, such as ketamine, play a role in preventing the development of central sensitization-a process that amplifies pain signals. Low-dose ketamine infusions have been used effectively to manage post-operative pain.¹² Alpha-2 agonists, like clonidine and dexmedetomidine, have analgesic properties and can be

used as adjuncts to anesthesia to improve pain control. They work by inhibiting norepinephrine release and dampening pain signals in the central nervous system.¹³

Patient-controlled analgesia is a technique that empowers patients to self-administer pre-programmed doses of intravenous analgesics, usually opioids like morphine or hydromorphone, to manage their pain. This approach gives patients better control over their pain relief while minimizing the risk of over sedation and optimizing analgesic delivery. Gabapentin is used as an additional pain management tool after surgery to decrease pain and lessen the need for opioids.¹⁴ It is effective in treating neuropathic pain but may cause drowsiness. A previous study investigated gabapentin use in surgery for patients aged 65 and older. The study found that using gabapentin during surgery led to higher risks of delirium, new antipsychotic use, and pneumonia in older patients.¹⁵ While gabapentin can aid pain relief during surgery, its benefits and drawbacks should be thoroughly assessed before prescribing it.

Overall, multimodal analgesia offers more comprehensive pain relief by targeting multiple pain pathways simultaneously. The use of non-opioid analgesics significantly reduces the risk of opioid-related side effects, such as respiratory depression, nausea, and constipation. It allows for the creation of tailored pain management plans that consider individual patient factors, ensuring personalized care and faster recovery, which ultimately leads to improved patient satisfaction. Multimodal analgesia makes patients suitable candidates for same-day discharge and promotes early resumption of

normal activities, resulting in cost savings for both patients and healthcare facilities.

Regional anesthesia

Regional anesthesia, which includes techniques like peripheral nerve blocks, epidural, and spinal anesthesia, is a specialized approach that selectively numbs specific nerves or nerve plexuses to block pain signals from reaching the brain. One significant advantage of regional anesthesia is the potential for early recovery, as the effects of anesthetics wear off more quickly, allowing for early ambulation and a smoother transition to the post-operative phase. Moreover, it offers excellent post-operative pain relief, reducing the reliance on systemic opioids. Patients are more actively involved in their care, which fosters a sense of control and lowers the risk of post-operative confusion or nausea.¹⁶ However, regional anesthesia does come with its own set of risks, including nerve damage, infection, and systemic toxicity from local anesthetics.¹⁷ Careful monitoring and skilled administration are crucial to minimize these potential complications.

Peripheral nerve blocks are commonly used for procedures in the extremities, such as hand and foot surgery, as well as certain orthopedic procedures. Epidural anesthesia involves injecting a local anesthetic into the epidural space around the spinal cord. On the other hand, spinal anesthesia, also known as intrathecal anesthesia, targets the cerebrospinal fluid and involves injecting a local anesthetic directly into the subarachnoid space, resulting in anesthesia from the level of injection downward.

Both techniques effectively block nerve impulses from the lower half of the body. However, these techniques cause complete nerve blockade (sensory, motor, proprioception, and autonomic), which can impair lower limb function for up to 6-8 hours. Due to these effects, they are generally not suitable for ambulatory surgery.

Managing chronic pain

Managing chronic pain in post-surgical patients is a critical aspect of healthcare, as a significant number of individuals experience persistent pain following surgery. While acute pain typically subsides within 1-2 weeks, chronic post-surgical pain can affect 10-50% of patients, significantly impacting their quality of life.¹⁸ Werner and Kongsgaard established specific criteria for diagnosing chronic post-surgical pain, which include pain that either appears new or intensifies after surgery, persists for more than 3-6 months post-surgery and substantially affects the patient's overall well-being, localized in the surgical area or in the nerve branches associated with that area, may persist since surgery or return after a period of relief, and is essential to differentiate this pain from other potential sources, such as cancer, infection, radiation therapy, or pre-existing conditions predating the surgery.¹⁹

Chronic pain patients often present with intricate medical backgrounds and may be managing various conditions like osteoarthritis, neuropathic pain, fibromyalgia, or chronic low back pain.²⁰ Identifying such patients is crucial during preoperative assessments to create customized pain management plans and address

issues related to opioid tolerance or dependence. Prior to the surgical procedure, collaboration between the ambulatory anesthesia team and pain specialists is essential. Pain specialists can provide valuable insights into adjusting chronic pain medications, while the anesthesia team can offer feedback on the patient's response to perioperative pain management strategies. Adjustments to chronic pain medications and the incorporation of non-opioid pain management strategies can significantly enhance pain control during the perioperative period. Multimodal analgesia is particularly pertinent for chronic pain patients. Given that chronic pain patients may have developed opioid tolerance, implementing opioid-sparing techniques like ketamine infusions or alpha-2 agonists can be beneficial in achieving adequate pain relief.²¹

Safety and complications

Notably, the use of opioids for pain relief can lead to various side effects, including nausea, vomiting, constipation, sedation, and respiratory depression. Allergic reactions to medications like local anesthetics or NSAIDs are also possible, along with risks associated with certain regional anesthesia techniques and nerve blocks, such as bleeding, hematoma formation, and nerve injury. To prioritize patient safety, collaboration with various healthcare professionals, including pain specialists, pharmacists, and physical therapists, ensures a thorough understanding of the patient's medical history, allergies, and current medications, allowing for the selection of the most appropriate pain management approach.

In addition to ensuring safety, educating

patients about potential complications and what to expect post-surgery is crucial for managing expectations and empowering patients to report any concerning symptoms promptly. Tailoring pain management plans to each patient's needs helps minimize the risk of adverse effects and enhances pain relief. Utilizing multimodal analgesia and considering patient-specific factors contribute to achieving optimal outcomes. Regularly monitoring vital signs, pain levels, and side effects during the perioperative period allows for prompt intervention if complications arise. Adhering to evidence-based guidelines and best practices in pain management is essential to providing patients with the most effective and safest treatments. Implementing standardized pain management protocols promotes consistency in care delivery and reduces the risk of errors. Moreover, integrating technology, such as electronic health records and automated monitoring systems, aids in patient safety, documentation, and data analysis.

Limitations and challenges

Traditional pain management methods often provide only short-term relief, leading to discomfort and potential complications during the initial stages of recovery. The reliance on opioids for pain control poses risks of side effects, including nausea, constipation, drowsiness, and the potential for opioid dependence. Inadequate pain control can result in delayed recovery and increased hospital readmission rates, affecting patient satisfaction and healthcare costs. Moreover, the one-size-fits-all approach to pain management does not consider individual patient factors, such

as pain tolerance and medical history, leading to suboptimal outcomes for some patients. In Thailand, the popularity of same-day surgery remains limited due to several challenges. These include a lack of specialized centers for systematic care, insufficient personnel to maintain the system, and inadequate equipment.

Ambulatory anesthesia offers significant advantages, but it also faces specific challenges related to pain management techniques. The duration of pain relief achieved through conventional methods, like local anesthetics or oral medications, may not always align with the necessary recovery period, resulting in pain resurfacing before patients have sufficiently healed. Managing pain effectively without over-relying on opioids remains a crucial challenge due to their potential for abuse, dependence, and side effects like nausea, constipation, and respiratory depression. Furthermore, each patient's pain threshold and response to pain management techniques can differ significantly, necessitating personalized approaches to find the most suitable and effective pain relief plan for individual patients.²² Inadequate patient education on pain management techniques, including medication usage and potential side effects, can also lead to non-compliance and suboptimal pain relief outcomes. Addressing these limitations and challenges is crucial for enhancing the overall patient experience and ensuring successful pain management in ambulatory anesthesia settings. Over the years, medical researchers and anesthesia providers have developed innovative approaches and incorporated cutting-edge technology to address these challenges,

ultimately improving patient comfort, safety, and recovery in outpatient settings.

Technology in ambulatory anesthesia pain management

Accurate pain assessment is paramount for effective pain management in ambulatory anesthesia. Advancements in technology have led to the development of various tools and methods to enhance pain assessment and quantification in ambulatory patients. Traditional pain scales, such as the numeric rating scale (NRS) and visual analog scale (VAS), have been digitized for electronic use. Patients can now conveniently indicate their pain levels using touch-screen devices or mobile applications, enabling quick and precise pain assessment.²³ Wearable devices equipped with sensors offer real-time monitoring of physiological parameters indicative of pain, such as heart rate, respiratory rate, and skin conductance. These data help healthcare providers gain a deeper understanding of patients' pain experiences and respond accordingly.

Patient-reported outcomes (PROs) are standardized questionnaires completed by patients at regular intervals to assess their pain experiences, medication use, and functional status. Incorporating PROs into electronic health records empowers healthcare providers to track and address patients' pain-related concerns more effectively.²⁴ Patient-controlled analgesia pumps allow patients to self-administer pre-programmed doses of pain medication, typically opioids, via intravenous or epidural routes. This patient-centered approach empowers individuals

to manage their pain effectively while minimizing the risk of overmedication. Transdermal patches deliver medications, such as opioids or non-opioid analgesics, through the skin and into the bloodstream over an extended period, offering continuous and convenient pain relief without frequent dosing. Additionally, intranasal and sublingual routes provide rapid and effective pain relief, eliminating the need for invasive procedures like intravenous access. Liposomal bupivacaine is a notable long-acting local anesthetic administered as a single injection. It gradually releases the drug over an extended period, delivering prolonged pain relief for up to 72 hours after surgery.²⁵

Innovative techniques have significantly contributed to improving pain control in ambulatory settings. Virtual reality (VR) and distraction techniques, for instance, have shown promise in reducing pain and anxiety during ambulatory procedures. VR allows patients to immerse themselves in calming virtual environments, effectively reducing their perception of pain and anxiety.²⁶ Similarly, activities such as listening to calming music or engaging in guided imagery can distract patients from pain and reduce discomfort. Gamified distraction techniques, such as interactive games or virtual experiences, effectively engage patients and minimize their focus on pain sensations.

Future directions and further research

Research in pain management should focus on developing predictive models that use patient-specific factors to tailor individualized

pain management plans. This could involve considering genetic variations, pain perception assessments, and other personalized factors. Additionally, exploring the efficacy of non-pharmacological pain management techniques, such as mindfulness, acupuncture, virtual reality, and music therapy, can help reduce opioid use and enhance patient comfort. Advancements in drug delivery systems, such as nanoparticle-based approaches or novel formulations, offer the potential to optimize pain relief while minimizing systemic side effects.²⁷ Moreover, investigating long-term pain management solutions for chronic pain patients undergoing ambulatory procedures can improve post-discharge care and reduce readmissions. Integrating technology, such as wearable devices, mobile apps, and virtual reality, can enhance pain assessment, patient engagement, and overall pain management outcomes.

Conclusion

Ambulatory anesthesia pain management has evolved significantly, incorporating innovative approaches to optimize patient care and outcomes. Patient-centered care, evidence-based protocols, and continuous quality improvement will remain essential elements in ensuring patient safety and comfort during ambulatory procedures. Personalized medicine, non-pharmacological interventions, drug delivery systems, and telehealth are promising areas that hold the potential to revolutionize pain management practices in outpatient settings. Embracing these advancements and actively exploring new horizons in pain management

research will enable healthcare providers to elevate the quality of care and improve patient outcomes in ambulatory anesthesia.

References

1. Lee JH. Anesthesia for ambulatory surgery. *Korean J Anesthesiol.* 2017;70:398-406.
2. Sohn HM, Ryu JH. Monitored anesthesia care in and outside the operating room. *Korean J Anesthesiol.* 2016;69:319-26.
3. Pregler JL, Kapur PA. The development of ambulatory anesthesia and future challenges. *Anesthesiol Clin North Am.* 2003;21:207-28.
4. Zwolinski NM, Patel KS, Vadivelu N, Kodumudi G, Kaye AD. ERAS protocol options for perioperative pain management of substance use disorder in the ambulatory surgical setting. *Curr Pain Headache Rep.* 2023;27:65-79.
5. Cukierman DS, Cata JP, Gan TJ. Enhanced recovery protocols for ambulatory surgery. *Best Pract Res Clin Anaesthesiol.* 2023;37:285-303.
6. Williams AC, Craig KD. Updating the definition of pain. *Pain.* 2016;157:2420-3.
7. Kehlet H. Procedure-specific postoperative pain management. *Anesthesiol Clin North Am.* 2005;23:203-10.
8. O'Neill A, Lirk P. Multimodal analgesia. *Anesthesiol Clin.* 2022;40:455-68.
9. Amaechi O, Huffman MM, Featherstone K. Pharmacologic therapy for acute pain. *Am Fam Physician.* 2021;104:63-72.
10. Husband M, Mehta V. Cyclo-oxygenase-2 inhibitors. *Cont Educ Anaesth Crit Care Pain.* 2013;13:131-5.
11. Schug SA, Chandrasena C. Postoperative pain management following ambulatory anesthesia: challenges and solutions. *Ambul Anesth.* 2015;2:11-20.
12. Riccardi A, Guarino M, Serra S, et al. Narrative review: low-dose ketamine for pain management. *J Clin Med.* 2023;12:3256.
13. Giovannitti JA Jr, Thoms SM, Crawford JJ. Alpha-2

- adrenergic receptor agonists: a review of current clinical applications. *Anesth Prog.* 2015;62:31-8.
14. Han C, Kuang MJ, Ma JX, Ma XL. The efficacy of preoperative gabapentin in spinal surgery: a meta-analysis of randomized controlled trials. *Pain Physician.* 2017;20:649-61.
 15. Park CM, Inouye SK, Marcantonio ER, et al. Perioperative gabapentin use and in-hospital adverse clinical events among older adults after major surgery. *JAMA Intern Med.* 2022;182:1117-27.
 16. Shams D, Sachse K, Statzer N, Gupta RK. Regional anesthesia complications and contraindications. *Clin Sports Med.* 2022;41:329-43.
 17. Kamel I, Ahmed MF, Sethi A. Regional anesthesia for orthopedic procedures: what orthopedic surgeons need to know. *World J Orthop.* 2022;13:11-35.
 18. Fuller AM, Bharde S, Sikandar S. The mechanisms and management of persistent postsurgical pain. *Front Pain Res (Lausanne).* 2023;4:1154597.
 19. Werner MU, Kongsgaard UE. Defining persistent post-surgical pain: is an update required. *Br J Anaesth.* 2014;113:1-4.
 20. Treede RD, Rief W, Barke A, et al. Chronic pain as a symptom or a disease: the IASP classification of chronic pain for the international classification of diseases (ICD-11). *Pain.* 2019;160:19-27.
 21. Ghai B, Jafra A, Bhatia N, Chanana N, Bansal D, Mehta V. Opioid sparing strategies for perioperative pain management other than regional anaesthesia: a narrative review. *J Anaesthesiol Clin Pharmacol.* 2022;38:3-10.
 22. Haverfield MC, Giannitrapani K, Timko C, Lorenz K. Patient-centered pain management communication from the patient perspective. *J Gen Intern Med.* 2018;33:1374-80.
 23. Thiel B, Godfried MB, van Huizen EC, et al. Patient reported postoperative pain with a smartphone application: a proof of concept. *PLoS One.* 2020;15:e0232082.
 24. Wang XS, Gottumukkala V. Patient-reported outcomes: is this the missing link in patient-centered perioperative care? *Best Pract Res Clin Anaesthesiol.* 2021;35:565-73.
 25. Malik O, Kaye AD, Kaye A, Belani K, Urman RD. Emerging roles of liposomal bupivacaine in anesthesia practice. *J Anaesthesiol Clin Pharmacol.* 2017;33:151-6.
 26. Chuan A, Zhou JJ, Hou RM, Stevens CJ, Bogdanovych A. Virtual reality for acute and chronic pain management in adult patients: a narrative review. *Anaesthesia.* 2021;76:695-704.
 27. Phadke A, Amin P. A recent update on drug delivery systems for pain management. *J Pain Palliat Care Pharmacother.* 2021;35:175-214.