

Anesthesia Practitioners' Inappropriate Handling with Narcotics and Psychotropics

Nuthakun Posri¹, Nisara Tanphan¹, Varathip Thongdech², Phongthara Vichitvejpaisal³

¹ Division of Perioperative Nursing, Golden Jubilee Medical Center, Faculty of Medicine Siriraj Hospital, Mahidol University, Nakhon Pathom 73170, Thailand

² Division of Nursing Specialty, Golden Jubilee Medical Center, Faculty of Medicine Siriraj Hospital, Mahidol University, Nakhon Pathom 73170, Thailand

³ Department of Anesthesiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

Abstract

In the complex landscape of healthcare, the phenomenon of anesthesia practitioners mishandling narcotics and psychotropics has surfaced as a pressing concern. This article delves into the enigmatic unraveling of this issue, shedding light on the implications it poses for patient safety and professional integrity. Anesthesia practitioners, entrusted with administering potent substances critical to pain management and patient well-being, stand at the intersection of ethical dilemmas and critical responsibilities. The article examines key points, emphasizing that patient safety is at immediate risk due to impaired decision-making and administration errors resulting from inappropriate substance use. Collaboration emerges as a cornerstone for addressing this challenge, necessitating cooperation among practitioners, institutions, regulatory bodies, and professional associations. Preventive measures and educational initiatives are underscored as pivotal strategies to mitigate substance abuse risks within this professional group.

Keywords: anesthesia practitioners, narcotics, psychotropics, substance abuse

Key points:

- Anesthesia practitioners' inappropriate use of narcotics and psychotropics compromises patient safety, as impaired judgment and decision-making can lead to errors in drug administration and vital sign monitoring.
- Addressing this concern demands collaboration among practitioners, institutions, regulatory bodies, and professional associations.
- Proactive prevention programs, stress management training, and educational initiatives can mitigate the risk of substance abuse among anesthesia practitioners.

Introduction

Anesthesia practitioners play a crucial role in ensuring patient safety and comfort during surgical procedures. They are responsible for administering controlled substances such as narcotics (opioids) and psychotropics (sedatives, tranquilizers) to manage pain, anxiety, and induce unconsciousness during surgeries. The demanding and high-stress nature of their profession, including the challenging work environment, extended hours, irregular schedules, exposure to trauma, and the demanding nature of patient

care, combined with easy access to potent medications, can render anesthesia practitioners vulnerable to substance abuse. This susceptibility can lead to addiction and its associated adverse consequences.¹

The misuse of narcotics, especially opioids, among anesthesia practitioners is of particular concern due to their potency and the potential for overdose. These drugs are not only highly addictive but also carry a significant risk of respiratory depression and other adverse effects. Such risks not only jeopardize the well-being of practitioners but also compromise the safety of their patients. Similarly, psychotropics, which are employed to manage stress and anxiety, can be abused, resulting in cognitive impairment, impaired decision-making, and compromised patient care. The primary objective of this article is to comprehensively examine the issue of substance abuse within the anesthesia practitioner community. By delving into the underlying causes, identifying risk factors, and proposing potential interventions to address the hidden challenges faced by anesthesia practitioners. These initiatives are designed not only to safeguard the mental and physical well-being of anesthesia practitioners but also to uphold the standard of patient safety.

Understanding narcotics and psychotropics

Narcotics and psychotropic substances are classes of drugs that have significant effects on the central nervous system, altering perception, mood, consciousness, and behavior. These substances are categorized based on their effects and potential for abuse, and they are subject to strict regulations due to their potential for dependence and harmful consequences.² Narcotics, also known as opioids, are a class of

drugs derived from opium or synthetic compounds with similar effects.² They are primarily used for pain management, especially in acute and postoperative settings, as they provide effective relief from severe pain, and can induce a sense of euphoria and relaxation.³ They work by binding to opioid receptors in the brain and spinal cord, blocking pain signals.

Psychotropic substances are drugs that affect mental processes, mood, and behavior. They are commonly used to manage psychiatric conditions such as anxiety, depression, insomnia, and various mental health conditions. These substances can be further classified into several categories, including:

Depressants slow down the central nervous system, and induce relaxation and sedation, help reduce anxiety and promote sleep, which can be beneficial for patients with insomnia or anxiety disorders. Depressants, when taken in excess, can cause drowsiness, impaired coordination, and even overdose. Examples include benzodiazepines, barbiturates, and certain sleep medications.⁴

Stimulants increase alertness, energy, and focus. They are used to treat conditions like attention deficit hyperactivity disorder, enhancing their daily functioning and narcolepsy. Stimulants, when abused, can lead to heart problems, anxiety, and paranoia. Examples include amphetamines and methylphenidate.⁴

Hallucinogenic substances alter perception, thoughts, and feelings. Although less commonly used in medical settings, hallucinations have shown promise in treating certain mental health issues, such as depression and post-traumatic stress disorder, when administered under controlled conditions. Hallucinogens, due to their unpredictable effects, can induce dangerous behavior and psychological distress. Examples include lysergic acid diethylamide (LSD),

psilocybin (magic mushrooms), and 3,4-methylenedioxymethamphetamine (MDMA).^{5,6} Many countries have stringent regulations and laws in place to govern the production, distribution, and possession of narcotics. Conversely, regulations pertaining to psychotropic drugs exhibit significant variation.

Roles and responsibilities of anesthesia practitioners

Anesthesia practitioners hold the crucial responsibility of guaranteeing patient comfort, safety, and optimal physiological conditions throughout surgical procedures. This entails conducting comprehensive preoperative assessments to ascertain patients' medical histories, allergies, current medications, and other pertinent factors. These details are utilized to formulate a precise anesthetic plan, encompassing suitable drugs, dosages, and techniques that could influence anesthesia administration.

In addition, anesthesia personnel are accountable for administering anesthetics, including narcotics and psychotropics, with precision and control. This ensures the induction and maintenance of anesthesia during surgical procedures. They maintain a vigilant watch over patients for any adverse reactions, making necessary adjustments in drug administration to uphold stable vital signs, such as heart rate, blood pressure, oxygen saturation, and respiratory rate. They also possess the ability to make real-time modifications to preserve stable physiological parameters.

Safeguarding an open and secure airway for unconscious patients is a key responsibility of anesthesia practitioners. This involves techniques like intubation, ventilation, and other methods aimed at maintaining optimal oxygenation and ventilation. Throughout the

surgical process and the post-operative period, they expertly manage pain by administering appropriate analgesics, thereby ensuring a comfortable recovery process. In cases of complications, these practitioners are trained to respond rapidly and effectively to emergencies, efficiently stabilizing the patient's condition and upholding their safety. Consequently, anesthesia providers serve as a vital link between the surgical team and the patient's well-being. This facilitates a seamless and controlled experience for both parties, significantly contributing to the success of surgical interventions.

Anesthesia practitioners exercise responsible medication usage, including narcotics and psychotropics, meticulously adhering to established protocols and dosages while considering the individual characteristics and needs of each patient. Despite the demanding and stressful nature of their profession, it's imperative for them to resist any inclination to misuse medications for personal relief. Instead, they should adopt healthy stress management strategies and coping mechanisms to mitigate the challenges associated with their role, thereby minimizing the risk of turning to substances for solace.

Substance abuse among anesthesia practitioners

Misuse of narcotics and psychotropic substances, not only jeopardizes the health and safety of the individual but also impacts patient care.⁷ Anesthesia practitioners, if affected by substance misuse, may experience impaired judgment, compromised patient monitoring, and decreased performance, potentially leading to medical errors and adverse patient outcomes.⁸ The incidence of substance use disorders among anesthesia practitioners was 2.7 times higher than that of medical professionals in other fields.^{9,10}

In light of this concerning statistic, there is a growing awareness of the imperative need for comprehensive preventive measures, education, and support systems within the anesthesia community to effectively address substance abuse issues. The routine use of narcotics and psychotropics in patient care can sometimes blur the boundaries between professional and personal use for some practitioners. This normalization of drug use within the profession contributes to an elevated risk of misuse. The ready accessibility of these substances can tempt individuals to misuse them, given their knowledge and easy access.¹¹ The insufficient monitoring and accountability measures within healthcare settings can make it easier for individuals to engage in substance misuse without immediate consequences.

Patterns of substance abuse

Patterns of narcotic abuse encompass various behaviors, including the misuse and abuse of prescription opioids through higher doses, frequent use, or obtaining prescriptions from multiple sources. Individuals may also engage in risky behaviors, such as “doctor shopping” or obtaining drugs from illicit sources, to sustain their substance use.¹² Additionally, narcotic abusers may combine these substances with other drugs, such as alcohol or benzodiazepines, to enhance their effects. Patterns of psychotropics abuse involve the unauthorized use of drugs like benzodiazepines and stimulants for their calming or stimulating effects, even without a prescription.¹³ Substances like MDMA (ecstasy) and LSD are commonly consumed for their hallucinogenic or empathogenic properties, often in social or recreational contexts.¹⁴ Psychotropics may be misused as a means of self-medicating to manage stress, anxiety, or depression without the oversight of medical professionals.

Substance abuse often leads to a cycle of tolerance, dependence, and escalating consumption as individuals chase the initial euphoric effects. Reducing the dose or abruptly stopping the drug may result in withdrawal symptoms and uncontrollable pain. This can involve increasing the dose to overcome tolerance, using the substance more frequently to avoid withdrawal symptoms, and transitioning to more potent administration methods, such as moving from oral consumption to injection.¹⁵ Additionally, mixing substances is common to intensify the high, but it heightens the risk of overdose and adverse reactions. An overdose involving both narcotics and psychotropics can induce sedation and suppress breathing, which can be fatal. The death rate has been found to be 10 times higher when combining narcotics and psychotropics compared to using narcotics alone.¹⁶

Psychological and environmental factors

Psychological factors encompass various elements, including stress, which is notably prevalent within the medical field and can compel individuals to turn to substances as a means of coping. Additionally, specific personality traits, such as impulsivity and a propensity for risk-taking, can contribute to the onset of substance abuse.¹⁷ Furthermore, underlying mental health issues, such as depression and anxiety, may also play a role in this context. Given the high stress levels commonly experienced in the medical profession, individuals may resort to unhealthy coping mechanisms, such as substance use, to alleviate emotional strain.¹⁸

Environmental factors include the demanding work environment in which anesthesia practitioners operate, which can lead to emotional strain and an increased likelihood of substance misuse.¹⁸ Peer influence and the culture within the workplace can normalize

substance use, making it easier for individuals to justify their actions. The isolated nature of working in the operating room can hinder colleagues from recognizing early signs of substance abuse and providing support. Furthermore, the accessibility of controlled substances in the workplace, coupled with lax monitoring, can tempt practitioners into misusing these drugs.¹¹

Factors affecting work environment

The unique stressors and dynamics within the work environment of anesthesia practitioners contribute to their vulnerability.¹¹ Working extensively alone in operating rooms, irregular and prolonged hours disrupt sleep and personal lives. The demanding nature of the profession, dealing with high-stakes situations, pursuing perfectionism, and the fear of errors, combined with pressure to ensure patient safety, lead to stress and burnout.⁸ These factors can escalate stress and potentially push some practitioners towards substance misuse as a coping mechanism, particularly when dealing with personal or stress-related challenges.^{11,18}

Moreover, isolation can hinder early detection of substance misuse and limit colleagues' ability to provide support. The stigma associated with seeking help for substance abuse, especially in esteemed fields like anesthesia, can deter individuals from seeking the needed assistance and treatment. Recognizing these challenges is crucial for implementing targeted interventions, raising awareness, and establishing support systems to address these concerns and uphold patient safety.

Impacts on patient care and safety

Substance abuse among anesthesia practitioners poses severe risks to patient care

and safety, undermining healthcare quality.^{16,17}

Cognitive impairment from substance abuse can impair judgment, leading to suboptimal decisions in critical moments of patient care, while diminished alertness and attention may hamper practitioners' ability to closely monitor vital signs and respond promptly to patient condition changes. Impaired coordination and motor skills can result in drug administration errors, including underdosing, overdosing, or administering the wrong medication. Slowed reflexes and coordination also hinder timely responses to emergencies, potentially causing delays in interventions and harm to patients.

Substance abuse can result in a multitude of errors and negligence that put patient care at risk, as impaired practitioners may administer incorrect dosages or the wrong medications, potentially causing inadequate pain relief, adverse reactions, or complications. The compromised ability to manage a patient's airway can lead to respiratory distress or failure, and impaired attention and reduced vigilance may result in inadequate monitoring of vital signs, potentially leading to the oversight of signs of deterioration. Impaired judgment may delay the recognition of critical situations, such as a drop in oxygen saturation or blood pressure, jeopardizing patient safety.

Additionally, substance abuse gives rise to substantial ethical concerns.³ Patients place their trust in these professionals to prioritize their well-being, but substance abuse undermines the practitioner's capacity to fulfill this trust, infringing upon patient autonomy and the principle of beneficence. Substance abuse may lead to dishonesty about one's condition or abilities, eroding the integrity of the practitioner-patient relationship. Practitioners grappling with substance abuse might prioritize personal needs over patient care, resulting in conflicts of interest and ethical dilemmas. Moreover,

substance abuse can impede transparency regarding errors or mistakes, obstructing the learning and improvement processes within healthcare systems.

Ethical and legal considerations

Healthcare professionals, including anesthesia practitioners, raise significant ethical concerns.³ These professionals have a primary ethical duty to ensure patient safety and well-being, but substance abuse compromises this responsibility, violating the principle of non-maleficence.¹¹ The obligation to be honest and transparent with patients is compromised when practitioners conceal their substance abuse or its effects, eroding trust and undermining the principle of veracity. Substance abuse can also result in professional misconduct and breaches of integrity, damaging the reputation of the healthcare profession. It may create a conflict between the practitioner's loyalty to their own needs and well-being and their commitment to patient care, thereby challenging the ethical principle of fidelity.

Healthcare professionals involved in substance abuse may encounter severe legal repercussions, including the potential revocation or suspension of their professional licenses, rendering them ineligible to practice. In cases involving diversion or illicit drug use, practitioners can face criminal charges ranging from drug possession and distribution to manslaughter if patient harm or death occurs. Errors or negligence resulting from substance abuse can expose practitioners to malpractice lawsuits, entailing legal claims for damages, while substance abuse-related mistakes can result in civil liability

cases, leading to financial penalties and harm to their professional reputation.

Healthcare institutions play a pivotal role in preventing and addressing substance abuse among anesthesia practitioners. To achieve this, institutions should establish educational programs that raise awareness about substance abuse, its consequences, and available resources for assistance. Routine monitoring of practitioners, drug inventories, and adherence to protocols can help identify early signs of substance abuse, with reporting mechanisms in place to facilitate timely intervention. Institutions should cultivate a supportive culture where practitioners feel safe seeking help for substance abuse without fear of reprisal. Furthermore, institutions should provide access to appropriate treatment and rehabilitation programs for those struggling with substance abuse, enabling their recovery and successful reintegration into practice.^{10,19}

Intervention, prevention and rehabilitation




Figure 1 3 key processes diagram

Table 1 Concepts of intervention, prevention, and rehabilitation

Intervention ^{10,19}	Prevention ^{19,20}	Rehabilitation ^{10,11}
Guidelines can help - Implement routine drug testing and monitoring - Encourage reporting of concerning behavior - Increase supervision and oversight - Provide education and training	Preventing drug use include - Incorporating substance abuse awareness into education and training - Establishing peer support networks and mentorship programs - Providing resources and training in stress management - Clear policies on substance abuse	Guidelines include - Providing access to evidence-based treatment programs - Facilitating a smooth return to work - Implementing monitoring and follow-up - Fostering a supportive culture

Collaborative efforts and future directions

Addressing substance abuse among anesthesia practitioners necessitates a collaborative approach involving multiple stakeholders. Anesthesia practitioners themselves must prioritize patient safety and well-being by supporting colleagues, reporting concerning behavior, and seeking help when substance abuse issues arise. Institutions should establish supportive environments that encourage reporting, provide resources for prevention and rehabilitation, and enforce policies against substance abuse. Regulatory bodies play a crucial role in setting standards and guidelines for addressing substance abuse, which includes implementing routine drug testing and monitoring. Simultaneously, professional organizations can promote awareness, education, and research initiatives related to substance abuse within the anesthesia community.^{11,21} Improving the process of monitoring, reporting, and addressing substance abuse cases can enhance patient safety and practitioner well-being through regular and random drug testing and real-time monitoring of vital signs. Practitioners struggling

with substance abuse should be offered non-punitive support, including access to rehabilitation and counseling services.¹¹

Ongoing research and awareness initiatives are vital to addressing substance abuse among anesthesia practitioners by identifying risk factors, effective interventions, and long-term outcomes.^{3,19} Publicizing the prevalence, consequences, and available resources can reduce stigma and encourage practitioners to seek help, while continuous education about the risks of substance abuse, stress management, and healthy coping strategies should be integrated into medical training and professional development.⁸ Collecting and analyzing data on substance abuse cases can provide insights into trends, enabling the development of targeted prevention and intervention strategies.³

Conclusion

Substance abuse jeopardizes patient safety and care quality by impairing judgment, attention, and decision-making, potentially leading to errors and delayed interventions. It also raises ethical dilemmas, breaches professional integrity, and poses legal conse-

quences, undermining the credibility of the healthcare profession. Addressing substance abuse requires collaborative efforts involving practitioners, institutions, regulatory bodies, and professional associations to prevent, intervene, and rehabilitate affected individuals. Strengthening monitoring protocols, reporting mechanisms, and intervention strategies is essential to identify and address cases of substance abuse promptly and supportively.

References

1. Samuelson ST, Bryson EO. The impaired anesthesiologist: what you should know about substance abuse. *Can J Anaesth*. 2017;64:219-35.
2. Narcotics Control Division, Food and Drug Administration. Act to implement the narcotics code 2021: narcotics code and narcotics case procedure act. 2nd ed. Bangkok: Aksorn Graphic and Design. 2022;p11-27.
3. Zhao S, Chen F, Feng A, Han W, Zhang Y. Risk factors and prevention strategies for postoperative opioid abuse. *Pain Res Manag*. 2019;2019:7490801.
4. Hilliard J. Central nervous system depressants. 2023 [cited 2023 Nov 29]. Available from <https://www.addictioncenter.com/drugs/drug-classifications/central-nervous-system-depressants/>.
5. De Gregorio D, Aguilar-Valles A, Preller KH, et al. Hallucinogens in mental health: preclinical and clinical studies on LSD, psilocybin, MDMA, and ketamine. *J Neurosci*. 2021;41:891-900.
6. Lowe H, Toyang N, Steele B, et al. The therapeutic potential of psilocybin. *Molecules*. 2021;26:2948.
7. Warner DO, Berge K, Sun H, Harman A, Wang T. Substance use disorder in physicians after completion of training in anesthesiology in the United States from 1977 to 2013. *Anesthesiology*. 2020;133:342-9.
8. Bryson EO. The opioid epidemic and the current prevalence of substance use disorder in anesthesiologists. *Curr Opin Anaesthesiol*. 2018;31:388-92.
9. Skipper GE, Campbell MD, Dupont RL. Anesthesiologists with substance use disorders: a 5-year outcome study from 16 state physician health programs. *Anesth Analg*. 2009;109:891-6.
10. Lefebvre LG, Kaufmann IM. The identification and management of substance use disorders in anesthesiologists. *Can J Anaesth*. 2017;64:211-8.
11. Misra U, Gilvarry E, Marshall J, et al. Substance use disorder in the anaesthetist: guidelines from the association of anaesthetists: guidelines from the association of anaesthetists. *Anaesthesia*. 2022;77:691-9.
12. Biernikiewicz M, Taieb V, Toumi M. Characteristics of doctor-shoppers: a systematic literature review. *J Mark Access Health Policy*. 2019;7:1595953.
13. Sanabria E, Cuenca RE, Estes MA, Maldonado M. Benzodiazepines: their use either as essential medicines or as toxic substances. *Toxics*. 2021;9:25.
14. Holze F, Vizeli P, Müller F, et al. Distinct acute effects of LSD, MDMA, and D-amphetamine in healthy subjects. *Neuropsychopharmacol*. 2020;45:462-71.
15. Agnoli A, Xing G, Tancredi DJ, Magnan E, Jerant A, Fenton JJ. Association of dose tapering with overdose or mental health crisis among patients prescribed long-term opioids. *JAMA*. 2021;326:411-9.
16. Dasgupta N, Funk MJ, Proescholdbell S, Hirsch A, Ribisl KM, Marshall S. Cohort study of the impact of high-dose opioid analgesics on overdose mortality. *Pain Med*. 2016;17:85-98.
17. Valdes JA. The concept of reentry in the addicted anesthesia provider. *AANA J*. 2014;82:95-100.
18. Andrade GO, Dantas RAA. Work-related mental and behaviour disorders in anesthesiologists. *Rev Bras Anesthesiol*. 2015;65:504-10.
19. Schonwald G, Skipper GE, Smith DE, Earley PH. Anesthesiologists and substance use disorders. *Anesth Analg*. 2014;119:1007-10.
20. DeFord S, Bonom J, Durbin T. A review of literature on substance abuse among anaesthesia providers. *J Res Nurs*. 2019;24:587-600.
21. Forget P, Baldacchino A. Substance abuse by anaesthesiologists, shouldn't we do more? *Eur J Anaesthesiol*. 2021;38:682-3.