Review Article • บทฟื้นฟูวิชาการ

Hypercapnia-Induced Complications in Anesthesia: a Current Practice

Tuanlutfee Tuwanhayee, Wilaiporn Supan, Kanittha Chongwilaiwan, Sirijanya Junyatham, Phongthara Vichitvejpaisal

Department of Anesthesiology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok 10700, Thailand

Abstract

Hypercapnia-induced complications in anesthesia are a critical concern for patient safety and surgical outcomes. This article provides a comprehensive analysis of current practices, physiological underpinnings, prevention, management, and future directions in the context of hypercapnia during anesthesia. It underscores the importance of vigilant carbon dioxide monitoring, timely intervention, and adherence to standardized protocols in mitigating the risk of hypercapnia. The physiological basis of hypercapnia, its induction during anesthesia, and its effects on various body systems are explained, emphasizing the significance of carbon dioxide homeostasis in acid-base balance and oxygenation. Pulmonary complications, cardiovascular effects, and neurological sequelae arising from hypercapnia are discussed, highlighting the potential impact on surgical outcomes and patient safety. The article reviews current monitoring and management practices, encompassing techniques, technologies, and ventilation strategies, with a focus on capnography. It emphasizes the need for patient-specific approaches, interdisciplinary communication, and adherence to standardized protocols. The discussion delves into patient selection, preoperative optimization, and anesthesia provider training. The article also explores future directions and innovations in carbon dioxide monitoring, equipment, and research areas requiring further exploration. Addressing hypercapnia in anesthesia is crucial for patient safety and surgical success. The article concludes by reiterating the importance of continuous education, customized care, interdisciplinary collaboration, and the incorporation of emerging technologies in improving patient care and outcomes.

Keywords: anesthesia, capnography, CO_2 monitoring, hypercapnia, ventilation

Introduction

Hypercapnia, defined as elevated levels of carbon dioxide (CO_2) in the bloodstream, is a critical consideration in the practice of anesthesia.[1] Hypercapnia can occur when there is an imbalance between CO_2 production, elimination, and regulation during the perioperative period, particularly when patients are under general anesthesia.[2] The potential for

Correspondence to: Phongthara Vichitvejpaisal, E-mail: phongthara@gmail.com

Received May 17, 2024, Revised April 23, 2025, Accepted April 23, 2025

hypercapnia poses significant challenges for anesthesia practitioners, as it can lead to a range of complications with the potential to impact patient safety and surgical outcomes. Continuous monitoring of CO_2 levels during surgery is now achievable through capnography. Hypercapnia can lead to respiratory and cardiovascular instability, posing a significant risk to patient safety. Furthermore, suboptimal CO_2 levels have the potential to affect tissue oxygenation and wound healing, underscoring the necessity to examine the link between hypercapnia and surgical outcomes. This article offers a comprehensive examination of hypercapnia within the realm of anesthesia, delving into its physiological underpinnings, causative factors, and its profound implications for patient well-being. Additionally, it discusses potential future innovations in anesthesia practice and offers insights into the evolving landscape of hypercapnia management.

Respiratory physiology

During normal respiration, the central chemoreceptor primarily detects changes in elevated CO_2 levels, while the peripheral chemoreceptor responds to reduced oxygen (O_2) tension, thereby adjusting the respiratory rate and depth accordingly. The lungs remove CO_2 from the bloodstream by exchanging it with O_2 in the alveoli. This gas exchange is crucial for maintaining adequate oxygenation and eliminating CO_2. CO_2 is generated as a byproduct of cellular metabolism, combining with water in the blood to form carbonic acid (H_2CO_3), which then dissociates into bicarbonate (HCO_3^-) and hydrogen ions (H+). Consequently, the most significant consequence of CO_2 is an impact on the body's acid-base balance. Elevated CO_2 levels result in an excess of H+, leading to respiratory acidosis. This disruption can compromise normal cellular function and affect various metabolic processes, such as impaired pulmonary function, altered cardiovascular dynamics, and changes in cerebral blood flow.[3,4]

Factors interrelated with intra-operative hypercapnia

Patient-specific factors

Patients with preexisting obstructive lung diseases, such as chronic obstructive pulmonary disease (COPD), obstructive sleep apnea (OSA), asthma, atelectasis (collapsed alveoli), ventilation-perfusion (V/Q) mismatch, as well as those who are pregnant, of advanced age, smokers, or obese, face an elevated risk due to compromised lung function, decreased lung compliance, and heightened airway resistance. Similarly, individuals with neuromuscular disorders, such as myasthenia gravis, are more susceptible to hypercapnia due to weakened respiratory muscles and a diminished ventilatory drive.[3,5,6]

Effective ventilation strategies

Anesthesia providers can exercise better control over minute ventilation and CO_2 levels by appropriately adjusting tidal volume and respiratory rate on the ventilator. Positive pressure ventilation can be employed to enhance lung recruitment and facilitate CO_2 elimination in patients with reduced lung compliance.[7] Additionally, a deliberate allowance of permissive hypercapnia, within safe limits, can minimize the risk of ventilator-induced lung injury. Continual capnography monitoring offers real-time feedback on exhaled CO_2 levels,

allowing providers to promptly fine-tune ventilation parameters in response to deviations.

Procedural aspects

Various factors during medical procedures can contribute to hypercapnia. Anesthetics, particularly sedative-hypnotics, opioids, benzodiazepines, and muscle relaxants, can depress a patient's ventilatory drive and decrease functional residual capacity.[8,9] Additionally, certain inhalational agents, including desflurane, sevoflurane, and isoflurane, may directly stimulate metabolism by promoting catecholamine release, particularly epinephrine and norepinephrine, consequently leading to heightened oxygen consumption and CO_2 production.

Physical airway obstructions, such as a dislodged or obstructed endotracheal tube, or bronchospasm, and patients with limited lung compliance, such as those undergoing surgeries necessitating a restrictive approach to lung ventilation, may experience CO_2 retention. Inappropriate settings on mechanical ventilators, such as insufficient tidal volume or excessive positive end-expiratory pressure (PEEP), malfunctioning ventilators or carbon dioxide absorbents, and specific surgical positions like steep Trendelenburg or lithotomy positioning, can result in V/Q mismatch, leading to inadequate CO_2 elimination. Certain surgical procedures, such as those utilizing CO_2 insufflation in laparoscopic surgeries or creating pneumoperitoneum, can introduce additional CO_2 into the body.[10] Furthermore, metabolic disorders such as diabetic ketoacidosis, severe metabolic acidosis, or sepsis, as well as hypermetabolic states induced by surgeries like thyroid or pheochromocytoma surgeries, can elevate CO_2 production.

Complications associated with hypercapnia

Pulmonary complications

Diminished alveolar ventilation, as observed in scenarios such as recurarization or inadequate mechanical ventilation, impacts CO_2 elimination, may lead to atelectasis, pneumonia, respiratory acidosis, and the exacerbation of preexisting lung conditions like COPD. These problems can result from reduced lung compliance and impaired mucus clearance due to bronchoconstriction. Prolonged hypercapnia can induce bronchoconstriction, heightened airway resistance, and fatigue of the respiratory muscles, ultimately causing breathing difficulties and potential respiratory distress. This may necessitate prolonged mechanical ventilation or the administration of supplemental oxygen.[11,12]

Cardiovascular effects

The cardiovascular system is highly responsive to fluctuations in CO_2 levels. Individual's physiological condition, the severity of hypercapnia, and the presence of other influencing factors or medical conditions are important.[13] Hypercapnia-induced vasodilation can result in a reduction in systemic vascular resistance, leading to hypotension.[3] It can also impact cardiac contractility and rhythm, potentially giving rise to arrhythmias or a decrease in cardiac output, which subsequently affects blood pressure and tissue perfusion. This is primarily because hypercapnia triggers the release of vasodilatory substances such as adenosine, nitric oxide, and prostaglandins. In conjunction with respiratory acidosis, these compounds directly relax the smooth muscles within the walls of blood vessels, further

promoting vasodilation. Conversely, hypercapnia may stimulate sympathetic activity, leading to the release of norepinephrine. This mechanism may serve as a protective response to sustain blood pressure by redirecting blood flow to vital organs. However, vasoconstriction can elevate heart rate and systemic vascular resistance, potentially resulting in hypertension and myocardial ischemia in individuals with preexisting cardiovascular conditions.[14,15]

Neurological sequelae

Hypercapnia can cause vasodilation in cerebral blood vessels, increasing intracranial pressure (ICP).[14] This can be especially concerning for patients with preexisting brain injuries, compromised cerebrovascular autoregulation, or those undergoing neurosurgical procedures. Elevated ICP may lead to cerebral edema and cerebral dysfunction, resulting in altered mental status, cognitive impairment, confusion, disorientation, or delirium in the postoperative period. Patients may experience difficulties with memory, attention, and problem-solving.[16,17]

Renal impairment

Prolonged hypercapnia may contribute to metabolic acidosis, electrolyte imbalances, and impaired renal perfusion. Hypercapnia-induced acidosis can adversely affect renal tubular function, electrolyte reabsorption, and fluid balance. Consequently, patients may develop hyponatremia, hyperkalemia, or volume depletion, predisposing them to renal dysfunction and acute kidney injury.[18]

Metabolic derangement

Increased CO_2 production during hypermetabolic states, such as major surgery, trauma, or sepsis, can exacerbate hypercapnia and metabolic acidosis. This escalation may heighten oxygen consumption, energy expenditure, and CO_2 generation, consequently further compromising respiratory and metabolic homeostasis. Elevated CO_2 levels can displace oxygen from hemoglobin molecules in the blood, reducing the oxygen-carrying capacity. Tissues and organs may receive insufficient oxygen, leading to tissue hypoxia.[19] This can result in respiratory acidosis, causing a decrease in blood pH and a more acidic environment. Respiratory acidosis can impair cellular function, potentially leading to arrhythmias, impaired oxygen release from hemoglobin, decreased cardiac contractility, impaired immune function, and delayed wound healing, rendering patients more susceptible to infections.[18]

Impact on patient outcomes

Complications arising from hypercapnia can prolong recovery times and necessitate extended stays in the post-anesthesia care unit or intensive care unit (ICU).[20] Insufficient tissue oxygenation can impair wound healing and increase the risk of nosocomial infections. Prolonged immobility and altered blood gas levels associated with hypercapnia can heighten the risk of deep vein thrombosis and pulmonary embolism. In severe cases, unmanaged hypercapnia, especially in patients with preexisting risk factors, can lead to life-threatening complications and increase the risk of perioperative mortality, particularly in patients with multiple comorbidities. These issues can extend hospital stays, delay discharges, raise healthcare costs, and decrease patient satisfaction.

Current CO_2 monitoring

Pulse oximetry primarily assesses oxygen saturation, but when combined with capnography, it can offer a more comprehensive view of respiratory status. Capnography, the gold standard for real-time CO_2 monitoring[21], particularly during anesthesia. It involves the continuous measurement of end-tidal CO_2 ($ETCO_2$) in exhaled breath, using infrared spectroscopy or mainstream/sidestream sampling. The capnogram provides insights into cardiac output and circulation, enhancing hemodynamic assessment. However, factors like water vapor interference, low cardiac output, or calibration issues can affect accuracy. Proper placement of sampling lines or cannulas is necessary.

In contrast, capnometry measures the partial pressure of CO_2 ($PetCO_2$) without graphical representation but provides numerical values correlating with arterial CO_2 levels. Although less comprehensive than capnography, it offers continuous feedback on ventilatory status, ensures correct endotracheal tube placement, and allows for immediate adjustments to ventilation and anesthetic management. Technical issues such as sampling line occlusion or disconnection may affect its reliability. End-tidal gas analysis monitors CO_2 concentration at expiration's end using gas chromatography or mass spectrometry, offering high accuracy and specificity, especially in situations where capnography or capnometry may be unreliable such as low tidal volumes, high respiratory rates, or low CO_2 levels. However, its expense and need for specialized equipment and expertise limit routine clinical use.

For non-invasive CO_2 monitoring, transcutaneous CO_2 monitoring is highly practical, providing continuous CO_2 level measurements through the skin using heated sensor electrodes. It benefits pediatric patients and those with compromised respiratory function, yet factors like skin integrity, temperature fluctuations, or electrode placement can affect accuracy and reliability. Calibration and periodic sensor replacement are necessary to maintain accuracy over time.[22] Esophageal CO_2 monitoring is employed in select cases, such as during cardiac surgeries or for patients with tracheostomies, to provide precise CO_2 monitoring. However, it's essential to note that arterial blood gas analysis, remains the most accurate method for assessing CO_2 levels in the bloodstream.[20]

Guidelines for managing hypercapnia

Anesthesia practitioners have several strategies at their disposal to effectively manage hypercapnia and maintain appropriate CO_2 levels in patients.

Tidal volume and respiratory rate adjustment

Anesthesia practitioners possess the expertise to meticulously adjust the tidal volume and respiratory rate parameters on ventilators. For instance, maintaining a tidal volume within the range of 6-8 mL/kg of ideal body weight and appropriately modifying the inspiratory time relative to the expiratory time (I:E ratio) from 1:2 to 1:1.5 optimizes CO_2 elimination during expiration. This approach

facilitates the achievement of normocapnia ($ETCO_2$ of 35-45 mmHg) and enhances ventilation efficiency.[23,24]

Positive pressure ventilation

The utilization of PPV stands as a valuable strategy to bolster lung recruitment and augment CO_2 elimination, particularly in patients grappling with compromised lung function. Nevertheless, exercising caution during configuration of settings is paramount to stave off complications such as barotrauma and excessive PEEP. It is imperative to customize PEEP levels in accordance with individual patient variables including lung compliance, airway resistance, and hemodynamic stability, with the goal of attaining the lowest effective PEEP to optimize respiratory mechanics. In some cases, it may be deliberately permissible to allow slightly elevated CO_2 levels, particularly in patients with severe lung disease. This approach, known as permissive hypercapnia, involves tolerating elevated CO_2 levels within safe limits.

Choice of mechanical ventilation mode

The choice between pressure-controlled ventilation (PCV) and volume-controlled ventilation (VCV) has a significant impact on CO_2 management. PCV may yield lower tidal volumes compared to VCV. The constant high pressure characteristic of PCV can help maintain open airways, making it suitable for patients with obstructive lung diseases or those at risk of airway collapse. VCV, on the other hand, ensures a consistent tidal volume with each breath, making it particularly suitable for patients with normal lung mechanics. It's essential to be mindful of patients with restrictive lung diseases, as VCV may lead to higher tidal volumes than necessary, potentially causing excessive CO_2 elimination and the risk of respiratory alkalosis.[25,26]

Hypercapnia prevention during anesthesia

A comprehensive preoperative assessment to identify patients at heightened risk of hypercapnia, such as those with preexisting COPD, asthma, OSA, smoking habits, obesity, and neuromuscular conditions is needed. Baseline respiratory function through spirometry, arterial blood gas analysis, and lung volume assessments to determine respiratory reserve are evaluated. Smoking cessation before surgery to mitigate the detrimental effects on lung function, implement respiratory therapy interventions including bronchodilators, mucolytics, and incentive spirometry, as well as optimize pulmonary function and enhance secretion clearance are done. In specific cases, optimize medication regimens aggressively, such as ensuring adherence to continuous positive airway pressure therapy for patients with OSA preoperatively, can alleviate respiratory symptoms and reduce perioperative complications.

Proper calibration of the mechanical ventilator and tailor ventilatory settings to individual patient needs, facilitate adequate CO_2 elimination.[27] Continuous monitoring of capnography and pulse oximetry enables prompt detection of deviations from normal CO_2 levels, such as esophageal intubation. Appropriate anesthetic depth, as indicated by the minimum alveolar concentration, and customize the reversal of muscle relaxants at the conclusion of surgery are maintained. For anesthesia

providers, standardized anesthesia protocols, endorsed by professional organizations, ensure anesthesia providers undergo regular comprehensive training and continuing education are implemented. They should collaborate with other healthcare professionals, including respiratory therapists, pulmonologists, and critical care specialists, to exchange expertise and best practices in managing respiratory complications.

Future directions

Emerging capnography developments include enhanced algorithms for more accurate monitoring, as well as miniaturized, wireless capnography devices that offer greater mobility and patient comfort. Closed-loop anesthesia systems, equipped with artificial intelligence and machine learning capabilities, have the potential to precisely adjust anesthetic agents, ventilation parameters, and CO_2 levels in real time.[28] Telemedicine and remote monitoring technologies are becoming more prevalent. Future ventilator modes may incorporate intelligent algorithms that adapt to a patient's respiratory patterns and needs, reducing the risk of hypercapnia. Anesthesia equipment is likely to become more integrated, with monitoring systems seamlessly interfacing with ventilators, anesthesia machines, and electronic health records. This integration can provide a comprehensive view of patient data in real time. The development of non-invasive CO_2 monitoring techniques, such as transcutaneous CO_2 monitoring, may become more precise and reliable, reducing the need for invasive arterial blood gas sampling.

Conclusion

The current practices in the management of hypercapnia during anesthesia underscore the need for continuous monitoring, vigilance, and a deep understanding of the physiological basis of hypercapnia. Elevated CO_2 levels can lead to a cascade of complications. Anesthesia practitioners should prioritize the observation of capnogram waveforms and be ready to intervene promptly when deviations occur. Patient-specific factors, including medical history and surgical requirements, should guide ventilation strategies. Teamwork ensures coordinated responses to hypercapnia-related issues. Emerging technologies enhance the prevention and management of hypercapnia.

References

1. Patel S, Miao JH. Hypercapnia. In: StatPearls. Treasure Island (FL): StatPearls Publishing; 2022[cited 2023 May 16]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK482456/
2. Foltz BD, Benumof JL. Mechanisms of hypoxemia and hypercapnia in the perioperative period. Crit Care Clin. 1987;3:269-86.
3. Curley G, Laffey JG, Kavanagh BP. Bench-to-bedside review: carbon dioxide. Crit Care. 2010;14:220.
4. Almanza-Hurtado A, Guerra CP, Martinez-Avila MC, Borre-Naranjo D, Rodriguez-Yanez T, Duenas-Castell C. Hypercapnia from physiology to practice. Int J Clin Pract. 2022;2022:2635616.
5. Malhotra A, Schwartz AR, Schneider H, et al. Research priorities in pathophysiology for sleep-disordered breathing in patients with chronic obstructive pulmonary disease. An Official American Thoracic Society Research Statement. Am J Respir Crit Care Med. 2018;197:289-99.
6. Karcz M, Papadakos PJ. Respiratory complications in the postanesthesia care unit: a review

of pathophysiological mechanisms. Can J Respir Ther. 2013;49:21-9.
7. Rittayamai N, Katsios CM, Beloncle F, Friedrich JO, Mancebo J, Brochard L. Pressure-controlled vs volume-controlled ventilation in acute respiratory failure: a physiology-based narrative and systematic review. Chest. 2015;148:340-55.
8. Boing S, Randerath WJ. Chronic hypoventilation syndromes and sleep-related hypoventilation. J Thorac Dis. 2015;7:1273-85.
9. Nimmo AF, Absalom AR, Bagshaw O, et al. Guidem lines for the safe practice of total intravenous anaesthesia (TIVA): Joint Guidelines from the Association of Anaesthetists and the Society for Intravenous Anaesthesia. Anaesthesia. 2019;74:211-24.
10. Sharma KC, Brandstetter RD, Brensilver JM, Jung LD. Cardiopulmonary physiology and pathophysp iology as a consequence of laparoscopic surgery. Chest. 1996;110:810-5.
11. Morales-Quinteros L, Camprubí-Rimblas M, Bringué J, Bos LD, Schultz MJ, Artigas A. The role of hypercapnia in acute respiratory failure. Intensive Care Med Exp. 2019;7(Suppl1):39.
12. Karcz M, Papadakos PJ. Respiratory complications in the postanesthesia care unit: a review of pathophysiological mechanisms. Can J Respir Ther. 2013;49:21-9.
13. Wilson MW, Labaki WW, Choi PJ. Mortality and healthcare use of patients with compensated hypercapnia. Ann Am Thorac Soc. 2021;18:e41
14. Curley G, Kavanagh BP, Laffey JG. Hypocapnia and the injured brain: more harm than benefit. Crit Care Med. 2010;38:1348-59.
15. Steinback CD, Salzer D, Medeiros PJ, Kowalchuk J, Shoemaker JK. Hypercapnic vs. hypoxic control of cardiovascular, cardiovagal, and sympathetic function. Am J Physiol Regul Integr Comp Physiol. 2009;296:R402-10.
16. Klein AA, Meek T, Allcock E, et al. Recommendations for standards of monitoring during anaesthesia and recovery 2021: Guideline from the Association of Anaesthetists. Anaesthesia. 2021;76:1212-23.
17. Zhou Q, Cao B, Niu L, et al. Effects of permis, sive hypercapnia on transient global cerebral ischemia–reperfusion injury in rats. Anesthesiology. 2010;112:288-97.
18. Kraut JA, Madias NE. Metabolic acidosis: pathophysiology, diagnosis and management. Nat Rev Nephrol. 2010;6:274-85.
19. Solhpour A, Tajbakhsh A, Safari S, Movaffaghi M, Pourhoseingholi MA, Soltani F. Inadvertent severe hypercapnia during general anesthesia: drop-in oxygen saturation or electrocardiography changes; which one warns us earlier? Ain-Shams J Anesthesiol. 2022;14:9.
20. Masterson C, Horie S, McCarthy SD, et al. Hypercapnia in the critically ill: insights from the bench to the bedside. Interface Focus. 2021;11:20200032.
21. Wilks C, Foran P. Capnography monitoring in the post anaesthesia care unit. J Periop Nurs. 2021;34:e29-35.
22. Mostafa MF, Osman AM, Abdallah ARM, Thabet AM, Hassan SA. Non-invasive carbon dioxide monitoring during moderate sedation at different oxygen flow rates in patients undergoing endoscopic retrograde cholangio-pancreatography. Egypt J Anaesth. 2023;39:63-71.
23. Tiruvoipati R, Gupta S, Bailey M, et al. Management of hypercapnia in critically ill mechanically ventilated patients-a narrative review of literature. J Intensive Care Soc. 2020;21:1-7.
24. Wright GT, Ashworth L, Pettey S. Optimizing mechanical ventilation during general anesthesia. AANA J. 2020;88:149-57.
25. Nin N, Muriel A, Peñuelas O, et al. Severe hyperu capnia and outcome of mechanically ventilated patients with moderate or severe acute respiratory distress syndrome. Intensive Care Med. 2017;43:200-8.
26. Bolther M, Henriksen J, Holmberg MJ, et al. Ventilation strategies during general anesthesia for noncardiac surgery: a systematic review and meta-analysis. Anesth Analg. 2022;135:971-85.
27. Coppola S, Froio S, Chiumello D. Protective lung ventilation during general anesthesia: is there any evidence? Crit Care. 2014;18:210.
28. Seger C, Cannesson M. Recent advances in the technology of anesthesia. F1000Res. 2020;9(F1000 Faculty Rev):375.