



Vet Integr Sci

Veterinary Integrative Sciences

ISSN: 2629-9968 (online)

Website: www.vet.cmu.ac.th/cmvj



Research article

Anthelmintic efficacy of mango seed extract on porcine gastrointestinal nematodes

Sentina Rhea T. Reyno¹, Loveille Jun A. Gonzaga², Elsa A. Gonzaga^{1,*}

¹Department of Veterinary Paraclinical Sciences, College of Veterinary Medicine, University of Southern Mindanao, Kabacan, 9407 Cotabato, Philippines

²Department of Chemistry, College of Science and Mathematics, University of Southern Mindanao, Kabacan, 9407 Cotabato, Philippines

Abstract

The usual control for gastrointestinal parasites is the use of commercial anthelmintics. However, parasites are becoming more resistant due to the frequent and inappropriate use of these anthelmintics. As a result, alternatives for these anthelmintics are becoming increasingly widespread for deworming livestock, particularly those that come from natural sources. Thus, this study aimed to evaluate the efficacy of mango (var. Carabao) seed extract in reducing the egg per gram counts of common gastrointestinal nematodes in pigs and compare the efficacy with that of levamisole. Experimental animals were naturally infected pigs with *Trichuris* spp. and *Strongyloides* spp. being the most predominant. The animals were administered orally with a single dose of the mango seed extract at concentrations of 600, 700, and 800 mg per kg bodyweight. The extract, regardless of concentration, was able to decrease the epg counts. Within 14 days post-treatment, there was no significant difference in the efficacies in the administration of 800 mg extract per kg bodyweight and levamisole. This comparable efficacy was sustained until 28 days post-treatment. This significant *in vivo* anthelmintic activity may be attributed to the tannins and flavonoids present in the extract. These results indicate that the mango seed extract is effective in controlling and reducing the gastrointestinal nematodes in pigs and may have the potential to be further developed as an anthelmintic.

Keywords: Anthelmintic, Gastrointestinal nematodes, Mango seed extract, Pigs

Corresponding author: Elsa A. Gonzaga, Department of Veterinary Paraclinical Sciences, College of Veterinary Medicine, University of Southern Mindanao, Kabacan, 9407 Cotabato, Philippines. Email: elsagonzaga@usm.edu.ph.

Article history; received manuscript: 8 May 2022,
revised manuscript: 16 May 2022,
accepted manuscript: 10 June 2022,
published online: 20 June 2022

Academic editor; Korakot Nganvongpanit



Open Access Copyright: ©2022 Author (s). This is an open access article distributed under the term of the Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution, and reproduction in any medium or format, as long as you give appropriate credit to the original author (s) and the source.

INTRODUCTION

Backyard farms are prone to disease outbreaks mainly due to inadequate facilities or poor disease monitoring. In the Philippines, there is a substantial number of smallholders or backyard pig raisers where it is used for their own consumption or is an important source of household income as a sideline or secondary source of income (Alawneh et al., 2014). Disease prevention and control remains an ongoing problem due to the costs of medicine (Villar et al., 2002). This results to parasitism being a common problem in backyard farms.

In particular, parasitic infection due to gastrointestinal nematodes is of economic importance because it affects the appetite, reduces weight gain due to poor feed utilization. If not controlled, heavy infection will result in high mortality (Roepstorff et al., 2011). In the Philippines, ascarids, strongyles, trichurids, and hookworms are commonly reported (Lee et al., 2005; Paller et al., 2019). The main mode of control or treatment remains to be the routine use of different anthelmintic products. However, the repeated and improper use of these anthelmintics particularly in smallholders often result on some degree of anthelmintic resistance (Pettersson et al., 2021). An alternative to these commercial anthelmintics is to explore plants or plant extracts that have the potential to be natural anthelmintics.

Mango (*Mangifera indica*) seed or generally known as stone, is thrown away as a waste product after consuming the fruit. Proximate analysis revealed that it contained polyphenolic compounds, among which tannins and vanillin were in highest amounts. Other compounds in the seed kernel include flavonoids, gallic acid, coumarin, caffeic acid, mangiferin, ferulic acid, and cinnamic acid (Abdalla et al., 2007). These compounds are reported to exert anthelmintic activities against a broad range of nematodes (Spiegler et al., 2017). These indicate that the mango seed extract could possibly be used as an anthelmintic against common gastrointestinal nematodes in pigs.

Thus, this study aimed to assess the efficacy of mango (var. Carabao) seed extract against common gastrointestinal nematodes in pigs. Specifically, it aimed to evaluate the efficacy of the mango seed extract in reducing the eggs per gram counts of common gastrointestinal nematodes in pigs in comparison to levamisole, a commercial anthelmintic.

MATERIALS AND METHODS

Ethics Approval

The procedures used in the study were approved by the University of Southern Mindanao Institutional Animal Care and Use Committee MC-CVM-001687. The methods for research and animal handling are all in compliance with Republic Act 8485, The Animal Welfare Act.

Animals

A preliminary quantitative examination of fecal samples was performed using the modified McMaster counting technique (Hutchinson, 2009) in order to determine the degree of parasitism. Pigs with more than 500 eggs per gram (epg) counts were used as experimental animals identified by tag markings. Fifteen pigs naturally-infected with gastrointestinal nematodes served as experimental animals. The pigs, between ages three to four months regardless of sex and breed, were used to assess the efficacy of different doses of mango seed extract in reducing the epg counts of common gastrointestinal nematodes in pigs. The pigs were from the same backyard farm and the farm's management systems and welfare measures were adapted. All diets were free of anthelmintics and were prepared to meet the nutritional requirements of the pigs. After the study, the experimental animals were released and the backyard farmer was informed of the results for possible interventions.

Collection and Preparation of Extract

Mangoes of the Carabao variety were purchased from the market of Kabacan, Cotabato, Philippines. The seeds were washed and the kernels were removed manually. The kernels were cut into smaller pieces and sundried. A mechanical grinder was used to produce a powder from the dried kernels. For extraction, the ground mango kernel (400 g) was soaked in 95% ethanol (2.40 L) for 48 hours and filtered using an ashless filter paper. The solvent was evaporated in vacuo at 40°C and 200 rpm, leaving around 12 g of crude extract for a 3% yield. The collected extracts were stored in a sterile container.

Administration of Extract

The fifteen naturally-infected pigs were divided into five homogeneous groups of three pigs each using randomized complete block design. This small sample size was because the extract was evaluated in vivo for the first time in this study. The identified experimental pigs were marked according to their group and were weighed using a calibrated weighing scale to determine the live weight. The treatments, extracts and controls, were administered orally early in the morning. Pigs in Groups A-C were treated with a single dose of 600, 700, and 800 mg extract per kg bodyweight, respectively. Group D was given a single dose of levamisole, 8.5 mg per kg bodyweight, as the positive control while Group E was not subjected to any treatment as the negative control. There were no adverse effects observed after the administration of the treatments.

Parasitological Examination

Fecal samples (3-5 g) were collected pre-treatment (day 0) and 7, 14, 21, and 28 days post-treatment by restraining the pig using a snare loop and inserting the fingers with gloves directly into the rectum. Fecal samples were placed in labeled empty plastic cups with cover prior to analysis. The animal care staff was unaware of the treatments administered to the pigs. Fecal examination was done using the modified McMaster technique (Hutchinson, 2009) to determine the fecal egg per gram counts. The efficacy was calculated as the difference between the post-treatment epg and pre-treatment epg, divided by the pre-treatment epg multiplied by 100.

Statistical Analysis

Results are expressed as means \pm standard errors of the means. One-way Analysis of Variance (ANOVA) was used to assess the significant differences ($p < 0.05$) in mean efficacy among the groups. In addition, post-hoc multiple mean concentration using the Tukey honestly significant difference (HSD) test was used when a significant difference was present and to test the consistency of the effects for treatments and control.

RESULTS

The experimental animals were raised in backyard farms thus mixed infection was observed. Among the naturally-infected pigs, the most predominant helminths identified were *Trichuris* spp. and *Strongyloides* spp., and to a lesser extent *Ascaris* spp.

The epg reduction regardless of the dosage of mango seed extract is evident from the data in Table 1. Among the extract concentrations, the highest reduction was seen using 800 mg of mango seed extract per kg bodyweight of pigs. Statistical analysis revealed that there was a significant difference among the epg counts 7, 14, 21, and 28 days post-treatment. Within 14 days post-treatment, the epg counts administered with 800 mg mango seed extract per kg bodyweight was comparable to that of the commercial anthelmintic levamisole. It should be noted that the epg counts started to increase for all groups within 21 days post-treatment, which continued until day 28.

The efficacies of the groups are shown in Table 2. The maximum efficacy for Group A was at 75.35% by day 7, which then continually decreased until day 28. For Group B, the maximum efficacy was observed 14 days post-treatment at 84.90%. Group C consistently elicited the highest mean efficacy among the mango seed extract concentrations for the entire duration of

Table 1 The mean fecal egg per gram of naturally-infected pigs treated with mango seed extract.

Group	Days post-treatment				
	0	7	14	21	28
A (600 mg/kg)	642 \pm 33 ^a	158 \pm 8 ^a	175 \pm 14 ^a	225 \pm 14 ^a	267 \pm 22 ^a
B (700 mg/kg)	550 \pm 14 ^a	108 \pm 8 ^{ab}	83 \pm 8 ^b	125 \pm 14 ^{ab}	167 \pm 8 ^b
C (800 mg/kg)	583 \pm 44 ^a	75 \pm 0 ^b	67 \pm 8 ^{bc}	100 \pm 14 ^{ab}	133 \pm 30 ^b
D (levamisole)	633 \pm 36 ^a	8 \pm 8 ^c	8 \pm 8 ^c	58 \pm 8 ^b	108 \pm 8 ^b
E (untreated)	550 \pm 14 ^a	583 \pm 22 ^d	600 \pm 29 ^d	683 \pm 46 ^c	767 \pm 22 ^c

*Mean values in the same column followed by the same superscript are not significantly different ($p < 0.05$).

the study. Administering 800 mg extract per kg bodyweight resulted to the highest mean efficacy among extract concentrations at 88.57% within 14 days post-treatment. Statistical analysis revealed that this efficacy is comparable to that of levamisole (98.81%). There was no significant difference between the

efficacies of Groups C and D, highlighting the potential of the extract to elicit similar anthelmintic activity with that of the commercial anthelmintic. It was further observed that the efficacy increased with increased concentration of the mango seed extract, implying a dose-dependent response.

Table 2 The mean efficacy of mango seed extract in different groups.

Group	Days post-treatment			
	7	14	21	28
A (600 mg/kg)	75.35 ± 1.26 ^a	72.79 ± 1.21 ^a	64.95 ± 1.08 ^a	58.56 ± 1.57 ^a
B (700 mg/kg)	80.34 ± 1.07 ^a	84.90 ± 1.16 ^b	77.31 ± 2.35 ^{ab}	69.73 ± 0.94 ^{ab}
C (800 mg/kg)	86.99 ± 1.03 ^b	88.65 ± 0.73 ^{bc}	83.03 ± 1.23 ^b	77.59 ± 4.10 ^b
D (levamisole)	98.81 ± 1.19 ^c	98.81 ± 1.19 ^c	90.86 ± 0.81 ^b	82.77 ± 1.65 ^b
E (untreated)	-6.00 ± 1.35 ^d	-9.11 ± 4.57 ^d	-24.08 ± 6.11 ^c	-39.52 ± 4.57 ^c

*Mean values in the same column followed by the same superscript are not significantly different ($p < 0.05$).

DISCUSSION

This efficacy of administering increasing dosages of mango seed extract resulting on the reduction of epg counts may be attributed to the anthelmintic components present in the mango seeds. Mango seed kernel extracts have been shown to contain tannins, flavonoids, alkaloids, saponins, and glycosides (Kaur et al., 2010). Extracts have been reported to exhibit a wide range of activities such as antioxidant, antimicrobial (Torres-León et al., 2021), antifungal (Dorta et al., 2016), antidiarrheal (Yakubu and Salimon, 2015), antiproliferative against colon cancer cells (Ballesteros-Vivas et al., 2019), and tyrosinase inhibition (Maisuthisakul and Gordon, 2009).

The mango seed has been considered a good source of polyphenols. For Egyptian mango Hindi cultivar, a significant amount of phenolics and flavonoids are present in mango seed kernel extracts at 17,400 and 3,325 mg per 100 g seed (Abdel-Aty et al., 2018). Similarly, among three Chinese mango cultivars, the kernels contained 15.5 mg of gallotannins per g dry matter (Luo et al., 2014). Kernels of the Ubá variety from Brazil contained 82,540 mg per kg dry matter (Ribeiro et al., 2008). Across these studies, it was observed that the mango seed kernel contained the highest amounts of phenolics compared to the pulp and peel, making it a promising source of polyphenolics.

Gallates and gallotannins are the major family of compounds present in the mango seed kernel extract. Gallotannins are hydrolyzable tannins that contain gallic acid substituents (Sáyago-Ayerdi et al., 2013). When hydrolyzed by acids and under physiological conditions in the gastrointestinal tract, these gallotannins may release gallic acid (Tomás-Barberan et al., 2009). Most studies report the role of condensed tannins related to anthelmintic activity. However, hydrolyzable tannins are reported to exhibit similar anthelmintic activity with that of condensed tannins. Extracts containing only hydrolyzable tannins have similar activity with extracts which have both condensed and hydrolyzable tannins and can even have higher anthelmintic activity than extracts containing only condensed tannins (Katiki et al., 2013). Gallotannins isolated from *Camelia sinensis* have shown strong anthelmintic activity against

Caenorhabditis elegans (Mukai et al., 2008). Moreover, condensed tannins with gallic acid units had higher anthelmintic activity compared to that with no gallic acid units (Brunet and Hoste, 2006).

The anthelmintic activity of hydrolyzable tannins is mainly egg hatching and larval migration inhibition while condensed tannins only exert minor effects on egg hatching. This implies that hydrolyzable tannins have a different mode of action for its anthelmintic activity (Engström et al., 2016). It was shown that the size of the hydrolyzable tannins, overall flexibility, type and number of functional groups, and the linkages between the monomeric hydrolyzable tannins had different effects on the activity (Kiuchi et al., 1988; Min et al., 2015). Particularly, the molecular size and the number of free hydroxyl groups seem to be crucial elements for anthelmintic activity (Spiegler et al., 2017). This may be due to the difficulty of larger molecules to permeate into target cells or within the parasite. In contrast, smaller molecules will be able to move easily and bind with target molecules.

In addition to hydrolyzable tannins, mango seed kernel also contain a substantial amount of flavonoids, which may have also contributed towards the reduction of epg counts. It is hypothesized that flavonoids have a mode of action similar to that of tannins. This means that flavonoids may interfere with the biology of the nematodes through egg hatching or larval motility inhibition such as the activity shown by sainfoin against *Haemonchus contortus* (Barrau et al., 2005). Though in smaller amounts, the alkaloids present may also have anthelmintic effects. Alkaloids have been reported to inhibit egg hatching (da Silva et al., 2021) and may have the ability to intercalate with DNA synthesis of parasites (Maqbool et al., 2004).

Aside from the mango seed kernel, studies have shown that other parts of the mango have anthelmintic activity. Immature mango fruits have shown high in vivo efficacy yet moderate anthelmintic action against nematodes, particularly *H. contortus*, in sheep (Nery et al., 2012). However, it was not reported as to what may have caused the activity nor the decrease in activity when it was tested in vivo. In contrast, this study was able to exhibit a strong anthelmintic action in vivo. This is largely due to the difference of the metabolism between sheep and pig but may also be due to the amount of the active substance in different parts of the mango. This study shows that the mango seed contains significant amounts of bioactive compounds enough to have an efficacy similar to a commercial anthelmintic.

The animals used in the study were pigs that are naturally infected with gastrointestinal nematodes and had at least 500 egg per gram counts, limiting the experimental animals. Increasing the sample size of the experimental animals may give a clearer response to the administration of the extract. Moreover, as previous studies on mango fruit extract reported strong in vitro yet moderate in vivo anthelmintic activity in sheep, this study focused on the in vivo anthelmintic activity in pigs. Future studies may be done to target specific gastrointestinal nematodes in vitro and in vivo as this study already supports that there is strong activity in vivo for mixed infection in pigs.

CONCLUSION

The results of the study show that the mango seed extract has potential *in vivo* anthelmintic activity against gastrointestinal nematodes in pigs. Administering 800 mg of the extract per kg bodyweight is able to elicit a similar efficacy with that of the commercial anthelmintic levamisole within 14 days post-treatment. This suggests that the anthelmintic activity of the mango seed extract is enough to warrant the potential to be developed as a naturally-sourced anthelmintic.

ACKNOWLEDGEMENTS

The expertise of Dr. Elizabeth C. Molina, Dr. Josephine R. Flores, and Dr. Rolando J. Garduque are much appreciated for the improvement of the study. The Tuklas Lunas Development Center facility, led by Prof. Lydia C. Pascual, had been vital in the conduct of the study.

AUTHOR CONTRIBUTIONS

SRTR, LJAG, and EAG developed the research concepts and design. SRTR conducted the experiments under the supervision and guidance of EAG. LJAG wrote the manuscript draft. All authors read and approved the final manuscript.

CONFLICT OF INTEREST

The authors declare no conflicts of interests.

REFERENCES

- Abdalla, A.E., Darwish, S.M., Ayad, E.H., El-Hamahmy, R.M., 2007. Egyptian mango by-product 1. Compositional quality of mango seed kernel. *Food Chem.* 103(4), 1134-1140.
- Abdel-Aty, A.M., Salama, W.H., Hamed, M.B., Fahmy, A.S., Mohamed, S.A., 2018. Phenolic-antioxidant capacity of mango seed kernels: therapeutic effect against viper venoms. *Rev. Bras. Farmacogn.* 28(5), 594-601.
- Alawneh, J.I., Barnes, T.S., Parke, C., Lapuz, E., David, E., Basinang, V., Baluyut, A., Villar, E., Lopez, E.L., Blackall, P.J., 2014. Description of the pig production systems, biosecurity practices and herd health providers in two provinces with high swine density in the Philippines. *Prev. Vet. Med.* 114(2), 73-87.
- Ballesteros-Vivas, D., Alvarez-Rivera, G., García Ocampo, A.F., Morantes, S.J., del Pilar Sánchez Camargo, A., Cifuentes, A., Parada-Alfonso, F., Ibáñez, E., 2019. Supercritical antisolvent fractionation as a tool for enhancing antiproliferative activity of mango seed kernel extracts against colon cancer cells. *J. Supercrit. Fluids.* 152, 104563.
- Barrau, E., Fabre, N., Fouraste, I., Hoste, H., 2005. Effect of bioactive compounds from Sainfoin (*Onobrychis viciifolia* Scop.) on the *in vitro* larval migration of *Haemonchus contortus*: role of tannins and flavonol glycosides. *Parasitology.* 131(4), 531-538.

- Brunet, S., Hoste, H., 2006. Monomers of condensed tannins affect the larval exsheathment of parasitic nematodes of ruminants. *J. Agric. Food Chem.* 54(20), 7481-7487.
- da Silva, G.D., de Lima, H.G., de Sousa, N.B., de Jesus Genipapeiro, I.L., Uzêda, R.S., Branco, A., Costa, S.L., Batatinha, M.J.M., Botura, M.B., 2021. In vitro anthelmintic evaluation of three alkaloids against gastrointestinal nematodes of goats. *Vet. Parasitol.* 296, 109505.
- Dorta, E., González, M., Lobo, M.G., Laich, F., 2016. Antifungal activity of mango peel and seed extracts against clinically pathogenic and food spoilage yeasts. *Nat. Prod. Res.* 30(22), 2598-2604.
- Engström, M., Karonen, M., Ahern, J., Baert, N., Payré, B., Hoste, H., Salminen, J.P., 2016. Chemical structures of plant Hydrolyzable Tannins reveal their in vitro activity against egg hatching and motility of *Haemonchus contortus* nematodes. *J. Agric. Food Chem.* 64(4), 840-851.
- Hutchinson, G., 2009. Nematode parasites of small ruminants, camelids and cattle diagnosis with emphasis on anthelmintic efficacy and resistance testing. *Australia and New Zealand Standard Diagnostic Procedures*, Canberra, pp. 17-19.
- Katiki, L.M., Ferreira, J.F., Gonzalez, J.M., Zajac, A.M., Lindsay, D.S., Chagas, A.S., Amarante, A.F., 2013. Anthelmintic effect of plant extracts containing condensed and hydrolyzable tannins on *Caenorhabditis elegans*, and their antioxidant capacity. *Vet. Parasitol.* 192(1-3), 218-27.
- Kaur, J., Rathinam, X., Kasi, M., Leng, K., Ayyalu, R., Kathiresan, S., Subramaniam, S., 2010. Preliminary investigation on the antibacterial activity of mango (*Mangifera indica* L.: Anacardiaceae) seed kernel. *Asian Pac. J. Trop. Med.* 3(9), 707-710.
- Kiuchi, F., Nakamura, N., Tsuda, Y., Kondo, K., Yoshimura, H., 1988. Studies on crude drugs effective on visceral larva migrans. IV. Isolation and identification of larvicidal principles in pepper. *Chem. Pharm. Bull. (Tokyo)*. 36(7), 2452-2465.
- Lee, J.A., Lañada E.B., More, S.J., Cotiw-an, B.S., Taveros, A.A., 2005. A longitudinal study of growing pigs raised by smallholder farmers in the Philippines. *Prev. Vet. Med.* 70(1-2), 75-93.
- Luo, F., Fu, Y., Xiang, Y., Yan, S., Hu, G., Huang, X., Huang, G., Sun, C., Li, X., Chen, K., 2014. Identification and quantification of gallotannins in mango (*Mangifera indica* L.) kernel and peel and their antiproliferative activities. *J. Funct. Foods*. 8, 282-291.
- Maisuthisakul, P., Gordon, M.H., 2009. Antioxidant and tyrosinase inhibitory activity of mango seed kernel by product. *Food Chem.* 117(2), 332-341.
- Maqbool, A., Hayat, C., Tanveer, A., 2004. Comparative efficacy of various indigenous and allopathic drugs against fascioliasis in buffaloes. *Vet. Arh.* 74(2), 107-114.
- Min, B., Hernandez, K., Pinchak, W., Anderson, R., Miller, J., Valencia, E., 2015. Effects of plant tannin extracts supplementation on animal performance and gastrointestinal parasites infestation in steers grazing winter wheat. *Open J. Anim. Sci.* 5(3), 343-350.
- Mukai, D., Matsuda, N., Yoshioka, Y., Sato, M., Yamasaki, T., 2008. Potential anthelmintics: polyphenols from the tea plant (*Camellia sinensis* L.) are lethally toxic to *Caenorhabditis elegans*. *J. Nat. Med.* 62, 155-159.
- Nery, P.S., Nogueira, F.A., Oliveira, N.J., Martins, E.R., Duarte, E.R., 2012. Efficacy of extracts of immature mango on ovine gastrointestinal nematodes. *Parasitol. Res.* 111(6), 2467-2471.
- Paller, V.G.V., Babia-Abion, S., 2019. Soil-transmitted helminth (STH) eggs contaminating soils in selected organic and conventional farms in the Philippines. *Parasite Epidemiol. Control.* 7, e00119.
- Pettersson, E., Halvarsson, P., Sjölund, M., Grandi, G., Wallgren, P., Höglund, J., 2021. First report on reduced efficacy of ivermectin on *Oesophagostomum* spp. on Swedish pig farms. *Vet. Parasitol. Reg. Stud. Reports.* 25, 100598.
- Ribeiro, S., Barbosa, L., Queiroz, J., Knödler, M., Schieber, A., 2008. Phenolic compounds and antioxidant capacity of Brazilian mango (*Mangifera indica* L.) varieties. *Food Chem.* 110(3), 620-626.
- Roepstorff, A., Mejer, H., Nejsun, P., Thamsborg, S.M., 2011. Helminth parasites in pigs: new challenges in pig production and current research highlights. *Vet. Parasitol.* 180(1-2), 72-81.

- Sáyago-Ayerdi, S.G., Moreno-Hernández, C.L., Montalvo-González, E., García-Magaña, M.L., Mata-Montes de Oca, M., Torres, J.L., Pérez-Jiménez, J., 2013. Mexican 'Ataulfo' mango (*Mangifera indica* L) as a source of hydrolyzable tannins. analysis by MALDI-TOF/TOF MS. *Food Res. Int.* 51(1), 188-194.
- Spiegler, V., Liebau, E., Hensel, A., 2017. Medicinal plant extracts and plant-derived polyphenols with anthelmintic activity against intestinal nematodes. *Nat. Prod. Rep.* 34(6), 627-643.
- Tomás-Barberan, F.A., Espín, J.C., García-Conesa, M.T., 2009. Bioavailability and metabolism of ellagic acid and ellagitannins. In: Quideau, S. (Ed.), *Chemistry and biology of ellagitannins*, World Scientific Publishing Co. Pte. Ltd., Singapore, pp. 273-297.
- Torres-León, C., de Azevedo Ramos, B., dos Santos Correia, M.T., Carneiro-da-Cunha, M.G., Ramirez-Guzman, N., Alves, L.C., Brayner, F.A., Ascasio-Valdes, J., Álvarez-Pérez, O.B., Aguilar, C.N., 2021. Antioxidant and anti-staphylococcal activity of polyphenolic-rich extracts from Ataulfo mango seed. *LWT.* 148, 111653.
- Villar, E., Maleon, M., Facundo, D., 2002. The swine industry in the Philippines: a focus on smallholders. In: *priorities for pig research in Southeast Asia and the Pacific to 2010*. Australian Centre for International Agricultural Research, Canberra, pp. 48-56.
- Yakubu, M.T., Salimon, S.S., 2015. Antidiarrhoeal activity of aqueous extract of *Mangifera indica* L. leaves in female albino rats. *J. Ethnopharmacol.* 163, 135-141.

How to cite this article;

Sentina Rhea T. Reyno, Loveille Jun A. Gonzaga and Elsa A. Gonzaga. Anthelmintic efficacy of mango seed extract on porcine gastrointestinal nematodes *Veterinary Integrative Sciences*. 2022; 20(3): 507- 515.
